

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 43/04, C07H 21/04, C12N 15/09, 15/11, 15/85		A1	(11) International Publication Number: WO 99/40789
			(43) International Publication Date: 19 August 1999 (19.08.99)
(21) International Application Number: PCT/US98/17908		(74) Agents: HANSBURG, Daniel; Kimeragen, Inc., 300 Pheasant Run, Newtown, PA 18940 (US) et al.	
(22) International Filing Date: 28 August 1998 (28.08.98)			
(30) Priority Data: 60/074,497 12 February 1998 (12.02.98) US 09/108,006 30 June 1998 (30.06.98) US		(81) Designated States: AL, AM, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(71) Applicants (for all designated States except US): REGENTS OF THE UNIVERSITY OF MINNESOTA [US/US]; 1100 Washington Avenue South, Minneapolis, MN 55414-1226 (US). ALBERT EINSTEIN COLLEGE OF MEDICINE OF YESHIVA UNIVERSITY [US/US]; 1300 Morris Park Avenue, Bronx, NY 10461 (US).		Published With international search report.	
(72) Inventors; and (75) Inventors/Applicants (for US only): STEER, Clifford, J. [US/US]; 1924 Pinchurst Avenue, St. Paul, MN 55116-1339 (US). KREN, Betsy, T. [US/US]; 810 29th Avenue S., Minneapolis, MN 55454 (US). BANDYOPADHYAY, Paramita, T. [IN/US]; 23208 Cornerstone Drive, Yardely, PA 19067 (US). ROY-CHOWDHURY, Jayanta [US/US]; 139 Woodhollow Lane, New Rochelle, NY 10804-3435 (US).			

(54) Title: METHODS AND COMPOUNDS FOR THE GENETIC TREATMENT OF HYPERLIPIDEMIA

(57) Abstract

The present invention concerns the introduction of specific alterations in the genes that encode three apolipoproteins, Apo A1, Apo B and Apo E. The alternations in Apo A1 introduce a cysteine residue so the disulfide cross-linked Apo A1 homodimers and Apo A1/A2 heterodimers can be formed. The alterations in Apo B introduce stop codons or frame shift mutations that cause the production of a truncated Apo B protein. The alterations in Apo E introduce specific point mutations that have been identified as protective. The production in the liver of a subject of these altered proteins reduces the risk of the subjects developing atherosclerosis. In one embodiment the genetic alterations are introduced by use of chimeric, mixed RNA/DNA, duplex oligonucleotides.

ATTORNEY DOCKET NUMBER: 7991-086-99
SERIAL NUMBER: 09/685,403
REFERENCE: BN



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 43/04, C07H 21/04, C12N 15/09, 15/11, 15/85	A1	(11) International Publication Number: WO 99/40789 (43) International Publication Date: 19 August 1999 (19.08.99)
(21) International Application Number: PCT/US98/17908 (22) International Filing Date: 28 August 1998 (28.08.98) (30) Priority Data: 60/074,497 12 February 1998 (12.02.98) US 09/108,006 30 June 1998 (30.06.98) US (71) Applicants (for all designated States except US): REGENTS OF THE UNIVERSITY OF MINNESOTA [US/US]; 1100 Washington Avenue South, Minneapolis, MN 55414-1226 (US). ALBERT EINSTEIN COLLEGE OF MEDICINE OF YESHIVA UNIVERSITY [US/US]; 1300 Morris Park Avenue, Bronx, NY 10461 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): STEER, Clifford, J. [US/US]; 1924 Pinehurst Avenue, St. Paul, MN 55116-1339 (US). KREN, Betsy, T. [US/US]; 810 29th Avenue S., Minneapolis, MN 55454 (US). BANDYOPADHYAY, Paramita, T. [IN/US]; 23208 Cornerstone Drive, Yardely, PA 19067 (US). ROY-CHOWDHURY, Jayanta [US/US]; 139 Woodhollow Lane, New Rochelle, NY 10804-3435 (US).		(74) Agents: HANSBURG, Daniel; Kimeragen, Inc., 300 Pheasant Run, Newtown, PA 18940 (US) et al. (81) Designated States: AL, AM, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: METHODS AND COMPOUNDS FOR THE GENETIC TREATMENT OF HYPERLIPIDEMIA		
(57) Abstract <p>The present invention concerns the introduction of specific alterations in the genes that encode three apolipoproteins, Apo A1, Apo B and Apo E. The alternations in Apo A1 introduce a cysteine residue so the disulfide cross-linked Apo A1 homodimers and Apo A1/A2 heterodimers can be formed. The alterations in Apo B introduce stop codons or frame shift mutations that cause the production of a truncated Apo B protein. The alterations in Apo E introduce specific point mutations that have been identified as protective. The production in the liver of a subject of these altered proteins reduces the risk of the subjects developing atherosclerosis. In one embodiment the genetic alterations are introduced by use of chimeric, mixed RNA/DNA, duplex oligonucleotides.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

METHODS AND COMPOUNDS FOR THE GENETIC TREATMENT OF HYPERLIPIDEMIA

This application claims benefit of the priority of application Serial No. 60/074,497, filed February 12, 1998 and application Serial No. 09/108,006 filed June 30, 1998.

1. FIELD OF THE INVENTION

The invention concerns methods and compositions for the use of recombinagenic oligonucleobases *in vivo* for the correction of disease causing genetic defects and the prevention of disease by introducing genetic modifications into the genes that encode Apolipoprotein B (Apo B), Apolipoprotein E (Apo E) and Apolipoprotein A1 (Apo A1).

2. BACKGROUND TO THE INVENTION

2.1 THE USE OF CHIMERIC MUTATIONAL VECTORS TO EFFECT GENETIC CHANGES IN CULTURED CELLS

The inclusion of a publication or patent application in this specification is not an admission that the publication or the invention, if any, of the application occurred prior to the present invention or resulted from the conception of a person other than the present inventors.

The published examples of recombinagenic oligonucleobases are termed Chimeric Mutational Vectors (CMV) or chimeraplasts because they contain both 2'-O-modified ribonucleotides and deoxyribonucleotides.

An oligonucleotide having complementary deoxyribonucleotides and ribonucleotides and containing a sequence homologous to a fragment of the bacteriophage M13mp19, was described in Kmiec, E.B., et al., November 1994, Mol. and Cell. Biol. **14**, 7163-7172. The oligonucleotide had a single contiguous segment of ribonucleotides. Kmiec et al. showed that the oligonucleotide was a substrate for the REC2 homologous pairing enzyme from *Ustilago maydis*.

Patent publication WO 95/15972, published June 15, 1995, and counterpart U.S. Patent No. 5,565,350 (the '350 patent) described duplex CMV for the introduction of genetic changes in eukaryotic cells. Examples in a *Ustilago maydis* gene and in the

murine ras gene were reported. The latter example was designed to introduce a transforming mutation into the ras gene so that the successful mutation of the ras gene in NIH 3T3 cells would cause the growth in soft agar of a colony of cells ("transformation"). The '350 patent reported that the maximum rate of transformation of NIH 3T3 was less than 0.1 %, i.e., about 100 transformants per 10^6 cells exposed to the ras duplex CMV. In the *Ustilago maydis* system, the rate of transformants was about 600 per 10^6 . A chimeric vector designed to introduce a mutation into a human bcl-2 gene was described in Kmiec, E.B., February 1996, Seminars in Oncology **23**, 188.

A duplex CMV designed to repair the mutation in codon 12 of K-ras was described in Kmiec, E.B., December 1995, Advanced Drug Delivery Reviews **17**, 333-40. The duplex CMV was tested in Capan 2, a cell line derived from a human pancreatic adenocarcinoma, using LIPOFECTIN™ to introduce the duplex CMV into the Capan 2 cells. Twenty four hours after the duplex CMV was introduced, the cells were harvested and genomic DNA was extracted; a fragment containing codon 12 of K-ras was amplified by PCR and the rate of conversion estimated by hybridization with allele specific probes. The rate of repair was reported to be approximately 18%.

A duplex CMV designed to repair a mutation in the gene encoding liver/bone/kidney type alkaline phosphatase was reported in Yoon, K., et al., March 1996, Proc. Natl. Acad. Sci. **93**, 2071. The alkaline phosphatase gene was transiently introduced into CHO cells by a plasmid. Six hours later the duplex CMV was introduced. The plasmid was recovered at 24 hours after introduction of the duplex CMV and analyzed. The results showed that approximately 30 to 38% of the alkaline phosphatase genes were repaired by the duplex CMV.

WO 97/41411 and counterpart United States Patent No. 5,760,012 to E.B. Kmiec, A. Cole-Strauss and K. Yoon, and the publication Cole-Strauss, A., et al., September 1996, SCIENCE **273**, 1386 disclose duplex CMV that are used in the treatment of genetic diseases of hematopoietic cells, e.g., Sickle Cell Disease, Thalassemia and Gaucher Disease. United States Patent Application Serial No. 08/664,487, filed June 17, 1996, by E.B. Kmiec describes duplex CMV having non-natural nucleotides for use in specific, site-directed mutagenesis. The duplex CMV described in the applications and certain of the publications of Kmiec and his colleagues contain a central segment of

DNA:DNA homoduplex and flanking segments of RNA:DNA hybrid-duplex or 2'-OMe-RNA:DNA hybrid-duplex.

The work of Kmiec and his colleagues concerned cells that are mitotically active, i.e., proliferating cells, at the time they are exposed to CMV. Kmiec and colleagues used a CMV/liposomal macromolecular carrier complex in which the CMV were mixed with a pre-formed liposome or lipid vesicle. In such a complex the CMV are believed to adhere to the surface of the liposome.

Kren et al., June 1997, *Hepatology* **25**, 1462-1468, reported the successful use of a CMV in non-replicating, primary tissue-cultured rat hepatocytes to mutate the coagulation factor IX gene. Kren et al., March 1998, *Nature Medicine* **4**, 285 reported the use of a CMV *in vivo* to introduce a genetic defect in the same gene.

2.2 THE USE OF A POLYETHYLENIMINE MACROMOLECULAR CARRIER FOR *IN VIVO* AND *IN VITRO* TRANSFECTION

Branched chain polyethylenimine has been used as a carrier to introduce nucleic acids into eukaryotic cells both *in vivo* and *in vitro*. Boussif, O., et al., 1995, *Proc. Natl. Acad. Sci.* **92**, 7297; Abdallah, B. et al., 1996, *Human Gene Therapy* **7**, 1947. Boletta, A., et al., 1997, **8**, 1243-1251. The *in vitro* use of galactosylated polyethylenimine to introduce DNA into cultured HepG2 hepatocarcinoma cell lines is reported by Zanta, et al., October 1, 1997, *Bioconjugate Chemistry* **8**, 839-844. The coupling of a protein ligand, transferrin, to polyethylenimine and its use to introduce a test gene into cultured cells by use of the transferrin receptor is described in Kircheis, R., et al., 1997, *Gene Therapy* **4**, 409-418. Branched chain polyethylenimines contain secondary and tertiary amino groups having a broad range of pK's and, consequently these polyethylenimines have a substantial buffering capacity at a pH where polylysine has little or no capacity, i.e., less than about 8. Tang, M.K., & Szoka, F.C., 1997, *Gene Therapy* **4**, 823-832. The use of branched chain polyalkanylimines, including polyethylenimine as carriers for the introduction of nucleic acids into cells is described in WO 96/02655 to J-P. Behr et al.

The successful *in vivo* and *in vitro* use of linear polyethylenimine to transfect a gene is reported by Ferrari, S., et al., 1997, *Gene Therapy* **4**, 1100-1106. Compositions

comprising a linear polyalkanylimine and a nucleic acid as disclosed in patent publication WO 93/20090 to S. Stein et al.

2.3 THE USE OF A LIPOSOMAL CARRIER FOR *IN VIVO* TRANSFECTION

The use of liposomes or lipid vesicles to introduce DNA encoding a foreign protein into cells has been described. The most frequently used techniques adhere the DNA to the surface of a positively charged liposome, rather than encapsulating the DNA, although encapsulated DNA techniques were known. United States Patent Nos. 4,235,871 and 4,394,448 are relevant. The field is reviewed by Smith, J.G., et al., 1993, *Biochim. Biophys. Acta* **1154**, 327-340 and Staubinger, R.M., et al., 1987, *Methods in Enzymology* **185**, 512. The use of DOTAP, a cationic lipid in a liposome to transfect hepatic cells *in vivo* is described in Fabrega, A.J., et al., 1996, *Transplantation* **62**, 1866-1871. The use of cationic lipid-containing liposomes to transfect a variety of cells of adult mice is described in Zhu, N., et al., 1993, *Science* **261**, 209. The use of phosphatidylserine containing lipids to form DNA encapsulating liposomes for transfection is described in Fraley, R., et al., 1981, *Biochemistry* **20**, 6978-87.

2.4 THE USE OF THE ASIALOGLYCOPROTEIN RECEPTOR FOR HEPATOCELLULAR SPECIFIC TRANSFECTION

United States Patent Nos. 5,166,320 and 5,635,383 disclose the transfection of hepatocytes by forming a complex of a DNA, a polycationic macromolecular carrier and a ligand for the asialoglycoprotein receptor. In one embodiment, the macromolecular carrier was polylysine. The use of a lactosylcerebroside containing liposome to transfect a hepatocyte *in vivo* is described by Nandi, P.K., et al., 1986, *J. Biol. Chem.* **261**, 16722-16722. The use of asialofetuin-labeled liposomes to transfect liver cells with a reporter plasmid is described in Hara et al., 1995, *Gene Therapy* **2**, 764-788. The use of galactosylated polyethyleneimine to transfect cultured hepatocytes is described in Zanta M-A., et al. abst. pub. Oct. 1, 1997, *Bioconjugate Chem.*, **8**, 839-844.

2.5 APO B100, APO B48 AND THE REDUCTION OF SERUM LDL

Hepatic and Intestinal Lipoprotein Secretion: Both the liver and the intestines make and export lipoproteins for the transport of lipids. The lipoproteins are termed very low density lipoproteins (VLDL) and chylomicrons, respectively. VLDL and chylomicrons differ in size and in their major protein components. The major protein of VLDL is Apo B100, consisting of 4536 amino acids; the major protein of chylomicrons is Apo B48, which consists of the N-terminal 2152 amino acids of Apo B100. Apo B48 and Apo B100 are encoded by a single gene, the transcript of which is modified at nucleotide 6666 (codon 2179) by a sequence specific cytidine deaminase, termed apolipoprotein B mRNA editing enzyme (APOBE). The action of this enzyme converts a C to U and results in a stop codon.

Both VLDL, which contain Apo B100, and chylomicrons, which contain Apo B48 transport triglycerides in the vascular system to a delivery site. However, after triglyceride hydrolysis and delivery VLDL are transformed into LDL, while chylomicrons are not. High levels of circulating LDL *per se* and a high LDL:HDL ratio increase the risk of arterial atherosclerosis. Hence, it has been suggested that increasing the ratio of Apo B48 to Apo B100 would have a beneficial effect.

In many species of mammals, e.g., rats and mice, a high percentage of the lipid secretions of both liver and intestine contain Apo B48. Such species have markedly lower ratios of LDL:HDL. Greve J., et al., 1995, Proc. Zool. Soc., Calcutta, **47**, 93-100. In others, such as humans and rabbits, hepatocytes lack APOBE and the hepatocytes consequently produce only VLDL.

One strategy to reduce the atherosclerosis in humans has been to introduce the gene for the catalytic component of the apolipoprotein B editing enzyme (APOBEC-1) under the control of a constitutive promoter to convert Apo B100 transcripts into Apo B48 transcripts. The transient expression of APOBEC-1 in the hepatocytes of normal and genetically hyperlipidemic Watanabe rabbit does cause a transient reduction in the levels of LDL. Greeve, J., et al., 1996, J. Lipid Res. **37**, 2001-17. However, the uncontrolled production of APOBEC-1 is mutagenic and may cause hepatocellular hyperplasia and hepatocellular carcinoma. Yamanaka, S., et al., 1995, Proc. Natl. Acad. Sci. **92**, 8483-8487.

Individuals who are homozygous or mixed heterozygotes for genes encoding truncated Apo B100 have been observed. Malloy et al., 1981, J. Clin. Invest. **67**, 1441; Hardman, D.A., et al., 1991, J. Clin. Invest. **88**, 1722. These individuals have low or absent LDL. For example, deletion of nucleotides 5391-5394 results in a frame shift mutation and a shortened Apo B (B37). These patients are most often asymptomatic. Steinberg, D., et al., 1979, J. Clin. Invest. **64**, 292; Young, S.G., et al., 1988, Science **241**, 591; Young, S.G., 1987, J. Clin. Invest. **79**, 1831. Reviewed Linton, M.F., 1993, J. Lipid. Res. **34**, 521; Kane, J.P. & Havel, R.J., 1995, Chapt. 57, THE METABOLIC BASIS OF INHERITED DISEASE, ed. Scriver et al. (McGraw Hill, New York). Similarly, as many as 1 in every 3,000 persons has a serum cholesterol level of 100 mg/dl or less because the individual is heterozygous for a truncated Apo B gene. *Ibid.*, p. 1866.

Truncations that result in an Apo B that are shorter than Apo B 31 do not circulate. Truncated Apo B 86, 87 and 90 have been observed. Apo B 86 and Apo B 87, are not associated with LDL while Apo B 90 is. Each mutation is associated with hypobetalipoproteinemia. Linton, M.L., et al., 1990, Clin. Res. **38**, 286A (abstr.); Tennyson, G.E., et al., 1990, Clin. Res. **38**, 482A (abstr.); Kruhl, E.S., et al., 1989, Arteriosclerosis **9**, 856.

2.6 APO E POLYMORPHISM AND TYPE III HYPERLIPIDEMIA

Apolipoprotein E is the major ligand for the LDL receptor for lipoproteins that contain Apo B48. There are three allelic forms of human Apo E that differ from each other by one or two amino acids: Apo E2 (Cys¹¹² Cys¹⁵⁸); Apo E3 (Cys¹¹² Arg¹⁵⁸); and Apo E4 (Arg¹¹² Arg¹⁵⁸). There is considerable geographical variation in the prevalences of the alleles. Excluding Africa, E2 ranges between 4% and 12 %, E3 between 70% and 85% and E4 between 7.5 and 25%. In the Sudan, the prevalences are 8.1%, 61.9% and 29.1%, respectively. Mahley, R.W. & Rall, S.C., Jr., 1995, Chapt. 61, THE METABOLIC BASIS OF INHERITED DISEASE, ed. Scriver et al. (McGraw Hill, New York). Thus approximately 1% of the North American and European population are Apo E 2/2 homozygotes. Of these homozygotes approximately between 2% and 10% display type III hyperlipidemia. Paradoxically, however, Apo E 2/2 homozygotes that have not

developed overt Type III hyperlipidemia display lower than average LDL associated cholesterol. Davignon, J., 1988, *Arteriosclerosis* **8**, 1.

The E4 allele is also associated with increased incidence of a major disease, Alzheimer's Disease, and with increased risk of coronary artery disease. Roses, A.D., 1996, *Ann. NY Acad. Sci.* **802**, 50-57; Okumoto, K., & Fujiki, Y., 1997, *Nature Genetics* **17**, 263; Kuusi, T., et al., 1989, *Arteriosclerosis* **9**, 237. A polymorphism in the region 491 nt 5' to the transcription start site of the Apo E gene is also an independently associated with increased risk of Alzheimer's disease. Individuals homozygous for the -491-A genotype have an increased risk of Alzheimer's, while individuals homozygous or heterozygous for the -491 T genotype have no increased risk. Bullido, M.J., 1998, et al., *Nature Genetics* **18**, 69-71.

The E2 allele in most individuals is associated with the lowest levels of serum cholesterol and LDL. However, about 5% of E2/E2 homozygous persons who are subject to environmental or genetic stress develop type III hyperlipidemia. The most common stressors are hypothyroidism, untreated diabetes mellitus, alcoholism and marked weight gain. Removal of the stressor usually results in control of the hyperlipidemia. Rare patients with type III hyperlipidemia have mutant Apo E genes. Mahley & Rall, *ibid.* Table 61-5.

2.7

APO A1 AND HDL

High density lipoproteins (HDL) transport cholesterol and phospholipids from peripheral extrahepatic locations to the liver. In particular, HDL are believed to remove lipid deposits from vascular endothelial cells and that the observed negative correlation between the levels of HDL-cholesterol (HDL-C) and coronary artery disease is due to this function. Eisenberg, S., 1984, *J. Lipid Research* **25**, 1017; Gordon, D.J., et al., 1986, *Circulation* **74**, 1217. HDL are secreted by the liver and intestines as nascent HDL particles containing four molecules of apo A1, which is a 243 amino acid protein. The nascent HDL physically attract free cholesterol from cell membranes and/or other lipoproteins. The resulting particle contains apo A1, phospholipid, and cholesterol. Such particles are substrates for lecithin:cholesterol acyltransferase (LCAT), which esterifies the free cholesterol to cholesterol esters. The presence of the more hydrophobic cholesterol

esters transforms the nascent HDL initially to a more stable mature HDL3 and subsequently HDL3 particle.

Cholesterol ester transfer protein removes the esterified cholesterol from the HDL3 and HDL2 particles and transfers them into LDL and thence into hepatocytes through the LDL receptors.

Two mutations of apo A1 have been discovered wherein the levels of HDL are depressed. The mutations are missense mutations that replace an arginine with a cysteine amino acid. Weisgraber, K.H., et al., 1983, J. Biol. Chem. **258**, 2508 (apo A1 milano (Arg-Cys)¹⁷³); Bruckert, E., et al., 1997, Atherosclerosis **128**, 121 (apo A1 R151C (Arg-Cys)¹⁵¹). The mutation is dominant, i.e., HDL levels are depressed in affected individuals who are heterozygous for the mutation. An explanation for this is that the mutant apo A1 proteins form cystine linked heterodimers with wild apo A2 molecules, which are a minor apolipoprotein found in HDL, as well as homodimers.

Surprising even though the lipid profile, i.e., low HDL and elevated triglycerides, is one that ordinarily is associated with accelerated atherosclerosis, the mutations are not associated with atherosclerosis, but, rather, are believed to be protective of atherosclerosis.

The mechanism of protection is not established. A large (N = 33) retrospective epidemiologic study of apo A1 milano carriers shows that there is a low incidence of coronary artery disease in these individuals, however, the significance of this fact must be viewed in the context of similar low incidence among non-affected residents of the village Limone sul Garda. Gualandri, V., et al., 1985, Am. J. Hum. Gen. **37**, 1083. *In vitro* studies indicate that the mutant apo A1 is not more effective in recruiting plasma membrane cholesterol. Bielicki, J.K., et al., 1997, Arterioscler Thromb Vasc Biol **17**, 1637. There is direct evidence that the administration of exogenous apo A1 milano, but not wild type apo A1, in the form of a complex with phospholipids, is protective in an accepted model system of atherosclerosis. Ameli, S. et al., 1994, Circulation **90**, 1935; Shah, P.K., et al., 1998, Circulation **97**, 780.

3. SUMMARY OF THE INVENTION

The present invention concerns methods of treatment and/or prophylaxis which consists of the introduction of specific genetic alterations in genes of a subject individual. In one embodiment, the specific genetic alteration blocks the synthesis of Apo B100 and thereby reduces the level of LDL cholesterol. In an alternative embodiment, the specific alteration converts an Apo E4 allele to an Apo E3 or Apo E2 allele, which is associated with decreased risk of atherosclerosis and Alzheimer's Disease. In further alternative embodiments, the invention concerns the correction of inherited genetic defects in the genes of hepatocytes of individuals having a disease caused by such defects.

The present invention further comprises a method of treating and/or preventing atherosclerosis by causing mutations in the genes encoding apo A1 and compounds that are useful for the introduction of such mutations. The mutations useful for the practice of the invention are mutations that insert a cysteine residue that forms a cystine and results in the formation of homodimers and apo A2-containing heterodimers.

The embodiments of the invention can be practiced using any oligonucleotide or analog or derivative thereof, now known or hereafter developed, that can cause specific genetic alterations in the genome of the hepatocytes of the subject individual (hereafter a "recombinagenic oligonucleobase"), for example a chimeric mutational vector (CMV) as, for example, described in United States patent No. 5,565,350, No. 5,731,181, and No. 5,760,012. Alternatively, the recombinagenic oligonucleobase can be a heteroduplex mutational vector or a non-chimeric mutational vector as described in U.S. patent application No. 09/078,063 and No. 09/078,064, filed May 12, 1998, each of which are hereby incorporated by reference.

In a preferred embodiment the recombinagenic oligonucleobase is complexed with a macromolecular carrier to which is attached a specific ligand. The ligand is selected to bind to a cell-surface receptor that is internalized into hepatocytes through clathrin-coated pits into endosomes. The cell surface receptors that bind such ligands are termed herein "clathrin-coated pit receptors". Examples of hepatic clathrin-coated pit receptors include the low density lipoprotein (LDL) receptor and the asialoglycoprotein receptor.

In specific embodiments the macromolecular carrier can be 1) an aqueous-cored lipid vesicle of between 25 nm and 400 nm diameter, wherein the aqueous core contains the CMV; 2) a lipid nanosphere of between 25 nm and 400 nm diameter, having a lipid core, wherein the lipid core contains a lipophilic salt of the CMV; or 3) a polycationic salt of the CMV. Examples of polycations for such salts include polyethylenimine, polylysine and histone H1. In one embodiment the polycation is a linear polyethylenimine (PEI) salt having a mass average molecular weight greater than 500 daltons and less than 1.3 Md. Alternatively the polycation can be a branched-chain polyethylenimine.

4. BRIEF DESCRIPTION OF THE FIGURE

Figure 1 is a schematic of one embodiment of CMV useful in the invention.

Figures 2A-2C show the genomic sequence of human APO E gene with translation of exons. Introns are in lower case and exons are in upper case.

Figure 3A-3G shows the sequence of the human APO A1 gene (SEQ ID No. 59)

5. DEFINITIONS

The invention is to be understood in accordance with the following definitions.

An oligonucleobase is a polymer of nucleobases, which polymer can hybridize by Watson-Crick base pairing to a DNA having the complementary sequence.

Nucleobases comprise a base, which is a purine, pyrimidine, or a derivative or analog thereof. Nucleobases include peptide nucleobases, the subunits of peptide nucleic acids, and morpholine nucleobases as well as nucleosides and nucleotides.

Nucleosides are nucleobases that contain a pentosefuranosyl moiety, e.g., an optionally substituted riboside or 2'-deoxyriboside. Nucleosides can be linked by one of several linkage moieties, which may or may not contain a phosphorus. Nucleosides that are linked by unsubstituted phosphodiester linkages are termed nucleotides.

An oligonucleobase chain has a single 5' and 3' terminus, which are the ultimate nucleobases of the polymer. A particular oligonucleobase chain can contain nucleobases of all types. An oligonucleobase compound is a compound comprising one or more oligonucleobase chains that are complementary and hybridized by Watson-Crick base pairing. Nucleobases are either deoxyribo-type or ribo-type. Ribo-type nucleobases are

pentosefuranosyl containing nucleobases wherein the 2' carbon is a methylene substituted with a hydroxyl, alkyloxy or halogen. Deoxyribo-type nucleobases are nucleobases other than ribo-type nucleobases and include all nucleobases that do not contain a pentosefuranosyl moiety.

An oligonucleobase strand generically includes both oligonucleobase chains and segments or regions of oligonucleobase chains. An oligonucleobase strand has a 3' end and a 5' end. When a oligonucleobase strand is coextensive with a chain, the 3' and 5' ends of the strand are also 3' and 5' termini of the chain.

A region is a portion of an oligonucleobase, the sequence of which is derived from some particular source, e.g., a CMV having a region of at least 15 nucleotides having the sequence of a fragment of the human β -globin gene. A segment is a portion of a CMV having some characteristic structural feature. A given segment or a given region can contain both 2'-deoxynucleotides and ribonucleotides. However, a ribo-type segment or a 2'-deoxyribo-type segment contain only ribo-type and 2'-deoxyribo-type nucleobases, respectively.

6. DETAILED DESCRIPTION OF THE INVENTION

6.1 THE STRUCTURE OF THE CHIMERIC MUTATIONAL VECTOR

The Chimeric Mutational Vectors (CMV) are comprised of oligonucleobases, i.e., polymers of nucleobases, which polymers form Watson-Crick base pairs of purines and pyrimidines (hybridize), to DNA having the appropriate sequence. Each CMV is divided into a first and a second strand of at least 15 nucleobases each that are complementary to each other. The strands can be, but need not be, covalently linked. Nucleobases contain a base, which is either a purine or a pyrimidine or analog or derivative thereof. There are two types of nucleobases. Ribo-type nucleobases are ribonucleosides having a 2'-hydroxyl, substituted 2'-hydroxyl or 2'-halo-substituted ribose. All nucleobases other than ribo-type nucleobases are deoxyribo-type nucleobases. Thus, deoxy-type nucleobases include peptide nucleobases. As used herein, only a recombinagenic oligonucleobase that contains at least three contiguous ribo-type nucleobases that are hybridized to deoxyribo-type nucleobases are considered CMV.

The sequence of the first and second strands consists of at least two regions that are homologous to the target gene, i.e., have the same sequence as fragments of the target gene, and one or more regions (the "mutator regions") that differ from the target gene and introduce the genetic change into the target gene. The mutator region is located between homologous regions. In certain embodiments of the invention, each of the flanking homologous regions contains a ribo-type segment of at least three ribo-type nucleobases, that form a hybrid duplex, preferably at least six ribo-type nucleobases and more preferably at least ten ribo-type nucleobases in length, but not more than 25 and preferably not more than 20, more preferably not more than 15 ribo-type nucleobases. The hybrid-duplex-forming ribo-type oligonucleobase segments need not be adjacent to the mutator region. In certain embodiments of the invention the ribo-type oligonucleobase segments are separated from the mutator region by a portion of the homologous region comprising deoxyribo-type nucleobases. In these embodiments the mutator region is also composed of deoxyribo-type nucleobases. Accordingly, the mutator region and a portion of one or both homologous regions form an intervening segment of homo-duplex, which separates the two segments of hybrid-duplex.

The total length of all homologous regions is preferably at least 16 nucleobases and is more preferably from about 20 nucleobases to about 60 nucleobases in length.

Preferably, the mutator region consists of 20 or fewer bases, more preferably 6 or fewer bases and most preferably 3 or fewer bases. The mutator region can be of a length different than the length of the sequence that separates the regions of the target gene homology with the homologous regions of the CMV so that an insertion or deletion of the target gene results. When the CMV is used to introduce a deletion in the target gene there is no base identifiable as within the mutator region. Rather, the mutation is effected by the juxtaposition of the two homologous regions that are separated in the target gene. For the purposes of the invention, the length of the mutator region of a CMV that introduces a deletion in the target gene is deemed to be the length of the deletion. In one embodiment the mutator region is a deletion of from 6 to 1 bases or more preferably from 3 to 1 bases. Multiple separated mutations can be introduced by a single CMV, in which case there are multiple mutator regions in the same CMV. Alternatively multiple CMV can be used simultaneously to introduce multiple genetic changes in a single gene

or, alternatively to introduce genetic changes in multiple genes of the same cell. Herein the mutator region is also termed the heterologous region.

In one embodiment the CMV is a single oligonucleobase chain of between 40 and 100 nucleobases. In an alternative embodiment, the CMV comprises a first and a second oligonucleobase chain, each of between 20 and 100 bases; wherein the first chain comprises the first strand and the second chain comprises the second strand. The first and second chains can be linked covalently by other than nucleobases or, alternatively, can be associated only by Watson-Crick base pairings. In an alternative embodiment the CMV is a first strand which is a single oligonucleobase chain and a second strand, complementary to the first which consists of two oligonucleobase chains, which are linked to the first strand chain by linkers. The combined length of the two chains of the second strand is the length of the first strand.

Linkers: Covalent linkage of the first and second strands can be made by oligo-alkanediols such as polyethyleneglycol, poly-1,3-propanediol or poly-1,4-butanediol. The length of various linkers suitable for connecting two hybridized nucleic acid strands is understood by those skilled in the art. A polyethylene glycol linker having from six to three ethylene units and terminal phosphoryl moieties is suitable. Durand, M. et al., 1990, *Nucleic Acid Research* **18**, 6353; Ma, M. Y-X., et al., 1993, *Nucleic Acids Res.* **21**, 2585-2589. A preferred alternative linker is bis-phosphorylpropyl-trans-4,4'-stilbenedicarboxamide. Letsinger, R.L., et alia, 1994, *J. Am. Chem. Soc.* **116**, 811-812; Letsinger, R.L. et alia, 1995, *J. Am. Chem. Soc.* **117**, 7323-7328, which are hereby incorporated by reference. Such linkers can be inserted into the CMV using conventional solid phase synthesis. Alternatively, the strands of the CMV can be separately synthesized and then hybridized and the interstrand linkage formed using a thiophoryl-containing stilbenedicarboxamide as described in patent publication WO 97/05284, February 13, 1997, to Letsinger R.L. et alia.

In a further alternative embodiment the linker can be a single strand oligonucleobase comprised of nuclease resistant nucleobases, e.g., a 2'-O-methyl, 2'-O-allyl or 2'-F-ribonucleotides. The tetranucleotide sequences TTTT, UUUU and UUCG and the trinucleotide sequences TTT, UUU, or UCG are particularly preferred nucleotide linkers.

Nucleotides: In an alternative embodiment the invention can be practiced using CMV comprising deoxynucleotides or deoxynucleosides and 2'-O substituted ribonucleotides or ribonucleosides. Suitable substituents include the substituents taught by the Kmiec Application, C₁₋₆ alkane. Alternative substituents include the substituents taught by U.S. Patent No. 5,334,711 (Sproat) and the substituents taught by patent publications EP 629 387 and EP 679657 (collectively, the Martin Applications), which are hereby incorporated by reference. As used herein a 2' fluoro, chloro or bromo derivative of a ribonucleotide or a ribonucleotide having a substituted 2'-O as described in the Martin Applications or Sproat is termed a "2'-Substituted Ribonucleotide." Particular preferred embodiments of 2'-Substituted Ribonucleotides are 2'-fluoro, 2'-methoxy, 2'-propyloxy, 2'-allyloxy, 2'-hydroxylethyloxy, 2'-methoxyethyloxy, 2'-fluoropropyloxy and 2'-trifluoropropyloxy substituted ribonucleotides. In more preferred embodiments the 2'-Substituted Ribonucleotides are 2'-fluoro, 2'-methoxy, 2'-methoxyethyloxy, and 2'-allyloxy substituted nucleotides.

2'-Substituted Ribonucleosides are defined analogously. Particular preferred embodiments of 2'-Substituted Ribonucleosides are 2'-fluoro, 2'-methoxy, 2'-propyloxy, 2'-allyloxy, 2'-hydroxylethyloxy, 2'-methoxyethyloxy, 2'-fluoropropyloxy and 2'-trifluoropropyloxy substituted ribonucleotides. In more preferred embodiment on the 2'-Substituted Ribonucleosides are 2'-fluoro, 2'-methoxy, 2'-methoxyethyloxy, and 2'-allyloxy substituted nucleotides.

The term "nuclease resistant ribonucleoside" encompasses 2'-Substituted Ribonucleosides, including 2'-Substituted Ribonucleotides and also all 2'-hydroxyl ribonucleosides other than ribonucleotides. In a preferred embodiment, the CMV preferably includes at least three and more preferably six nuclease resistant ribonucleosides. In one preferred embodiment the CMV contains no nuclease sensitive ribonucleosides. In an alternative preferred embodiment, every other ribonucleoside is nuclease resistant. Certain 2'-blocking groups can be more readily synthesized for purines or pyrimidines. In one embodiment of the CMV only the ribonucleoside purines or only the ribonucleoside pyrimidines are nuclease resistant.

Recombinagenic oligonucleobases, including non-chimeric mutational oligonucleobases and improved CMV and their use in eukaryotic cells and cell-free

systems are described in U.S. patent applications Serial No. 09/078,063, filed May 12, 1998, and Serial No. 09/078,064, filed May 12, 1998, which are each hereby incorporated in their entirety. These mutational oligonucleobases can be used in the same manner as the CMV described in this application.

6.2 THE GENE-SPECIFIC STRUCTURE OF THE CHIMERIC MUTATIONAL VECTOR

Figure 1 shows a diagram of a CMV according to one embodiment of the invention. In the Figure segments "a" and "c-e" are target gene specific segments of the CMV. The sequence of segment "a" and "c-e" are complements of each other. The sequence of segments "f" and "h" are also complements of each other but are unrelated to the specific target gene and are selected merely to ensure the stability of hybridization in order to protect the 3' and 5' ends. Additional protection of the 3' and 5' ends can be accomplished by making the 5' and 3' most internucleotide bonds a phosphorothioate, phosphonate or any other nuclease resistant bond. The sequence of segments "f" and "h" can be 5'-GCGCG-3' or permutations thereof. Segments "g" and "b" can be any linker that covalently connects the two strands, e.g., four unpaired nucleotides or an alkoxy oligomer such as polyethylene glycol. When segments "g" and "b" are composed of other than nucleobases, then segments "a", "c-f" and "h" are each an oligonucleobase chain.

The ribo-type nucleobase segments are segments "c" and "e," which form hybrid-duplexes by Watson-Crick base pairing to the complementary portions of segment "a." The segment "a" can have the sequence of either the coding or non-coding strand of the gene.

Table I contains SEQ ID No. 4 - No. 21 and Table III contains SEQ ID No. 22-25 and 54-58, which are examples of the sequences that can be used to practice the invention. The mutator region in each case is underlined and in bold. CMV having a segment "a" with a sequence selected from the sequences of Table I can be used to practice the invention. Alternatively, segment "a" may have the sequence of the complement of a sequence of Table I. As used herein, a CMV or other type of recombinagenic oligonucleobase comprises a sequence if either strand of the CMV or recombinagenic oligonucleobase comprises the sequence or comprises a sequence containing ribo-type nucleobases with uracil bases replacing thymine bases. Thus, for

example, a CMV having the sequence 5'-agucuggaugGGTAAGccgcccua-3' (SEQ ID No. 26) is considered to have the sequence of SEQ ID No: 4, wherein the lower case letters denote ribo-type nucleobases and the UPPER CASE letters denote deoxyribo-type nucleobases.

Subjects can be treated with a recombinagenic oligonucleobase specific for Apo B or Apo E according to the guidance of the Factor IX example below. More particularly the recombinagenic oligonucleobase can be given in divided doses at intervals that permit determining of the phenotypic effect of the dose, i.e., evaluation of the extent of the decline in LDL cholesterol and observation for adverse reactions. A reduction of the subject's fasting LDL serum cholesterol to below the level of the 5th percentile of the age-matched population (80-90 mg/dl) can be used as a therapeutic end point; alternatively reduction of fasting LDL serum cholesterol to below the average age-matched normal value (100-140) can be used. The number and size of the dose(s) can be modified to control the extent of the phenotypic effects. In the event that reversal of the specific genetic changes appear desirable, a recombinagenic oligonucleobase having a sequence appropriate to reverse the specific changes can be administered so that the fraction of unmodified Apo B or Apo E genes can be increased. Modification of the dose size and number and the administration of a reversing recombinagenic oligonucleobase permits the adjustment of the number of altered genes in the subject so that a predetermined amount of the phenotypic change can be effected.

6.2.1 Specific Alterations of the Apo B Gene

SEQ ID No. 1 contains the Apo B amino acid sequence and SEQ ID No. 2 contains the Apo B cDNA sequence.

The level of serum cholesterol and particularly of LDL-associated cholesterol can be reduced in a subject by introducing mutations into the subject's hepatic Apo B genes. The mutation can be any mutation that causes termination of the Apo B translation product between amino acid 1433 (Apo B 31) and amino acid 3974 (Apo B 87). (The amino acid numbering for Apo B in this specification refers to the 4553 amino acid primary translation product, i.e., mature Apo B100 plus the 27 amino acid leader sequence. Mature Apo B 100 consists of 4536 amino acids and mature Apo B 48 consists

of 2152 amino acids.) Preferably the translation product is terminated between amino acids 1841 (Apo B 40) and 2975 (Apo 65). The translation product can be terminated by introducing a frameshift mutation, i.e., by adding or deleting one or two nucleotides from the gene, or by introducing a stop codon (a TAA, TAG or TGA). The preferred stop codon is TAA. To monitor the introduction of the mutation it is preferred to have the mutation introduce or remove a palindromic sequence, which is the substrate of a restriction enzyme.

The sequence of the CMV is selected to have two homologous regions of at least 10 nucleobases and preferably at least 12 nucleobases each with a fragment of the Apo B gene located between nucleotides encoding amino acid 1433 (nt 4425) and 3974 (nt 12,048) and preferably located between the nucleotides encoding amino acids 1841 (nt 5649) and 2975 (nt 9051). In this specification, nt 6666 is the first nucleotide of codon 2180, the nucleotide that is converted by APOBE. In a preferred embodiment, the two homologous regions are separated by a single nucleobase in the sequence of the Apo B gene, where the CMV introduces a base substitution in the Apo B gene. Alternatively, the two homology regions can be adjacent in the Apo B gene and separated by a single or double nucleobase in the CMV, such that a one or two base insertion results from the action of the CMV on the Apo B gene. Alternatively, the homologous regions can be separated in the Apo B gene by one or two nucleotides that are deleted from the sequence of the CMV, such that the action of the CMV results in a one or two base deletion in the gene.

Nucleotides 4425-12,048 of the Apo B cDNA are encoded by exon 26 (nt 4342-11913), exon 27 (nt 11914 - 12028) and exon 28 (nt 12029-12212); see Table I, and GENBANK Accession No. 19828, which is hereby incorporated by reference. When an alteration is to be made at a position 3' of nt 11913, attention must be paid to the exon/intron boundary. Mutations that are located within 10-15 nucleotides of the exon/intron boundary must be identified so that the homology region of the CMV continues with the sequence of the intron and not the exon.

The homologous regions can be each from 10 to about 15 nucleobases in length; the two regions need not be of the same length. The fraction of nucleobases that contain a guanine or cytosine base is a design consideration (the GC fraction). It is preferred that

when the homologous region contains 12 or fewer nucleobases, the GC fraction be at least 33% and preferably at least 50%. When the GC fraction is less than 33% the length of the homologous regions is preferably 13, 14 or 15 nucleobases.

Table I contains 18 exemplary embodiments, SEQ ID No. 4-21 and Table III contains 9 exemplary embodiments, SEQ ID No. 22-25 and 54-58, of CMV sufficient for the practice of the embodiments of the invention described in this section. Suitable CMV can be made using nt 3-23 of SEQ ID No. 4-10, 12, and 16-20. SEQ ID NO. 11 and 13-15 have a lower GC fraction; CMV sufficient for the practice of the invention can be made containing residues 3-25 of SEQ ID NO. 11 and 13-15.

6.2.2 Specific Alterations of the Apo E Gene

In a further embodiment, the invention consists of introducing specific alterations to the Apo E gene. E4 homozygous individuals are at increased risk for atherosclerosis, particularly coronary artery disease, and Alzheimer's disease. Therefore, one embodiment of the present invention is the introduction of the substitution Arg→Cys at residues 112, to convert an E4 allele to an E3 allele, and optionally at residue 158 to convert an E3 or E4 allele into an E2 allele of an Apo E gene of an hepatocyte of a subject. The substitutions can be introduced using an oligonucleobase containing the sequence of nt 3-23 of SEQ ID No. 22 and No. 23 or complement thereof and more preferably of an oligonucleobase containing SEQ ID No. 22 and No. 23 or complement thereof. In addition, in individuals lacking genetic or environment stressors, the E2 allele results in a lowered LDL level and a decreased risk of atherosclerosis and coronary artery disease. Thus, these risks in an E3/E3 individual can be reduced by introduction of the (Arg→Cys)¹⁵⁸ substitution to convert the individual Apo E genes to E2 alleles.

Apo E2/E2 homozygous individuals who are suffering from Type III hyperlipidemia can be treated by converting E2 alleles to E3 alleles by making a Cys→Arg¹⁵⁸ substitution. Such a substitution can be made using an oligonucleobase containing the sequence of nt 3-23 of SEQ ID No. 24 or complement thereof and more preferably of an oligonucleobase containing SEQ ID No. 24 or complement thereof.

Independent of the Apo E allele, individuals who are homozygous for -491-A are at increased risk to develop Alzheimer's Disease. Bullido, M.J., 1998, et al., Nature

Genetics 18, 69-71. These individuals can be advantageously treated with an oligonucleobase containing the sequence of nt 3-23 of SEQ ID No. 25.

6.2.3 Repair of Mutations of the Apo B and Apo E Gene

SEQ ID No. 3 contains the Apo E genomic DNA sequence.

A further embodiment of the invention concerns the use of CMV to repair mutations in the Apo B and Apo E genes that cause hypobetalipoproteinemia and dysbetalipoproteinemia, respectively. Mutations that are located within 10-15 nucleotides of the exon/intron boundary must be identified so that the homology region of the CMV continues with the sequence of the intron and not the exon. The genomic sequence of Apo E4 indicating the exon and intron boundaries is given in Paik et al., 1985, Proc. Natl. Acad. Sci. 82, 3445, which is hereby incorporated by reference. The exon/intron boundaries of the Apo B gene are given in Table II along with the GENBANK accession numbers for the genomic sequence of Apo B.

6.2.4 Specific Alterations of the Apo A1 Gene

Apo A1 is a 243 amino acid protein. Amino acids 99-230 are encoded by six tandem duplications of a 66 base pair prototype sequence. The duplications are between 80 % and 64 % homologous to the consensus sequence. Without limitation as to theory the conformation of amino acids 120 to 230 is believed to be helical. The sequence of the amino acids is such that the helix is amphipathic, i.e., the helix has a hydrophobic face and a hydrophilic face, which contains polar amino acids.

The mutations most suitable for the practice of the invention are substitutions of polar amino acids by cysteine. Particularly suitable mutations are at arginine residues that are located next to other polar amino acids because the arginine codon used in apo A1 (CGC) can be converted into a TGC cysteine codon by a single base change. Thus, particularly suitable sites for mutations according to the present invention are arginine 149, 151, 153, 171 and 173.


6.3 FORMULATIONS SUITABLE FOR *IN VIVO* USE

The prior art formulations of CMV and a macromolecular carrier are of limited utility for *in vivo* use because of their low capacity for CMV and because the CMV is not protected from extracellular enzymes. The invention provides three alternative macromolecular carriers that overcome the limitations of the prior art. The carriers are polyethylenimine (PEI), aqueous-cored lipid vesicles, which are also termed unilamellar liposomes and lipid nanospheres.

Each of the carriers can be further provided with a ligand that is complementary to a cell-surface protein of the target cell. Such ligands are useful to increase both the amount and specificity of the uptake of CMV into the targeted cell. In one embodiment of the invention the target cell is a hepatocyte and the ligand is a galactose saccharide or lactose disaccharide that binds to the asialoglycoprotein receptor.

6.3.1 Polycationic Carriers

The invention can be practiced using any polycation that is non-toxic when administered to cells *in vitro* or to subjects *in vivo*. Suitable examples include polybasic amino acids such as polylysine, polyarginine, basic proteins such as histone H1, and synthetic polymers such as the branched-chain polyethylenimine:



The invention can be practiced with any branched chain polyethylenimine (PEI) having an average molecular weight of greater than about 500 daltons, preferably greater than between about 10 Kd and more preferably about 25 Kd (mass average molecular weight determined by light scattering). The upper limit of suitability is determined by the toxicity and solubility of the PEI. Toxicity and insolubility of molecular weights greater than about 1.3 Md makes such PEI material less suitable. The use of high molecular weight PEI as a carrier to transfect a cell with DNA is described in Boussif, O. et al., 1995, Proc. Natl. Acad. Sci. **92**, 7297, which is hereby incorporated by reference. PEI solutions can be prepared according to the procedure of Boussif et al.

The CMV carrier complex is formed by mixing an aqueous solution of CMV and a neutral aqueous solution of PEI at a ratio of between 9 and 4 PEI nitrogens per CMV phosphate. In a preferred embodiment the ratio is 6. The complex can be formed, for

example, by mixing a 10 mM solution of PEI, at pH 7.0 in 0.15 M NaCl with CMV to form a final CMV concentration of between 100 and 500 nM.

In addition a ligand for a clathrin-coated pit receptor can be attached to the polycation or to a fraction of the polycations. In one embodiment the ligand is a saccharide or disaccharide that binds to the asialoglycoprotein receptor, such as lactose, galactose, or N-acetylgalactosamine. Any technique can be used to attach the ligands. The optimal ratio of ligand to polyethylene subunit can be determined by fluorescently labeling the CMV and injecting fluorescent CMV/molecular carrier/ligand complexes directly into the tissue of interest and determining the extent of fluorescent uptake according to the method of Kren et al., 1997, *Hepatology* 25, 1462-1468.

Good results can be obtained using a 1:1 mixture of lactosylated PEI having a ratio of 0.4-0.8 lactosyl moieties per nitrogen and unmodified PEI. The mixture is used in a ratio of between 4 and 9 PEI nitrogens per CMV phosphate. A preferred ratio of oligonucleotide phosphate to nitrogen is 1:6. Good results can be obtained with PEIs having a mass average molecular weight of 25 Kd and 800 Kd which are commercially available from Aldrich Chemical Co., Catalog No. 40,872-7 and 18,197-8, respectively. Linear PEI such as that described in Ferrarri, S., et al., 1997, *Gene Therapy* 4, 1100-1106 and sold under the trademark EXGEN 500™ is particularly suitable for the practice of the invention because of its lower toxicity compared to branched-chain PEI.

In an alternative embodiment the polycationic carrier can be a basic protein such as histone H1, which can be substituted with a ligand for a clathrin-coated pit receptor. A 1:1 (w/w) mixture of histone and CMV can be used to practice the invention.

6.3.2 Lipids that Are Useful in Carriers

The selection of lipids for incorporation into the lipid vesicle and lipid nanosphere carriers of the invention is not critical. Lipid nanospheres can be constructed using semi-purified lipid biological preparations, e.g., soybean oil (Sigma Chem. Co.) and egg phosphatidyl choline (EPC) (Avanti Polar Lipids). Other lipids that are useful in the preparation of lipid nanospheres and/or lipid vesicles include neutral lipids, e.g., dioleoyl phosphatidylcholine (DOPC), and dioleoyl phosphatidyl ethanolamine (DOPE), anionic lipids, e.g., dioleoyl phosphatidyl serine (DOPS) and cationic lipids, e.g., dioleoyl

trimethyl ammonium propane (DOTAP), dioctadecyldiamidoglycyl spermine (DOGS), dioleoyl trimethyl ammonium (DOTMA) and DOSPER (1,3-di-oleoyloxy-2-(6-carboxy-spermyl)-propyl-amide tetraacetate, commercially available from Boehringer-Mannheim). Additional examples of lipids that can be used in the invention can be found in Gao, X. and Huany, L., 1995, *Gene Therapy* 2, 710. Saccharide ligands can be added in the form of saccharide cerebroside, e.g., lactosylcerebroside or galactocerebroside (Avanti Polar Lipids).

The particular choice of lipid is not critical. Hydrogenated EPC or lysolecithin can be used in place of EPC. DPPC (dipalmitoyl phosphatidylcholine), can be incorporated to improve the efficacy and/or stability of the delivery system.

6.3.3 The Construction of Lipid Nanosphere Carriers

Lipid nanospheres can be constructed by the following process. A methanol or chloroform methanol solution of phospholipids is added to a small test tube and the solvent removed by a nitrogen stream to leave a lipid film. A lipophilic salt of CMV is formed by mixing an aqueous saline solution of CMV with an ethanolic solution of a cationic lipid. Good results can be obtained when the cationic species are in about a 4 fold molar excess relative to the CMV anions (phosphates). The lipophilic CMV salt solution is added to the lipid film, vortexed gently followed by the addition of an amount of neutral lipid equal in weight to the phospholipids. The concentration of CMV can be up to about 3% (w/w) of the total amount of lipid.

After addition of the neutral lipid, the emulsion is sonicated at 4°C for about 1 hour until the formation of a milky suspension with no obvious signs of separation. The suspension is extruded through polycarbonate filters until a final diameter of about 50 nm is achieved. When the target cell is a reticuloendothelial cell the preferred diameter of the lipid nanospheres is about 100-200 nm. The CMV-carrying lipid nanospheres can then be washed and placed into a pharmaceutically acceptable carrier or tissue culture medium. The capacity of lipid nanospheres is about 2.5 mg CMV/ 500 μ l of a nanosphere suspension.

6.3.4 The Construction of Lipid Vesicles

A lipid film is formed by placing a chloroform methanol solution of lipid in a tube and removing the solvent by a nitrogen stream. An aqueous saline solution of CMV is added such that the amount of CMV is between 20% and 50% (w/w) of the amount of lipid, and the amount of aqueous solvent is about 80% (w/w) of the amount of lipid in the final mixture. After gentle vortexing the liposome-containing liquid is forced through successively finer polycarbonate filter membranes until a final diameter of about 50 nm is achieved. The passage through the successively finer polycarbonate filter results in the conversion of polylamellar liposomes into unilamellar liposomes, i.e., vesicles. When the target cell is a reticuloendothelial cell the preferred diameter of the lipid nanospheres is about 100-200 nm. The CMV-carrying lipid nanospheres can then be washed and placed into a pharmaceutically acceptable carrier or tissue culture medium.

The CMV are entrapped in the aqueous core of the vesicles. About 50% of the added CMV is entrapped.

A variation of the basic procedure comprises the formation of an aqueous solution containing a PEI/CMV condensate at a ratio of about 4 PEI imines per CMV phosphate. The condensate can be particularly useful when the liposomes are positively charged, i.e., the lipid vesicle contains a concentration of cations of cationic lipids such as DOTAP, DOTMA or DOSPER, greater than the concentration of anions of anionic lipids such as DOPS. The capacity of lipid vesicles is about 150 μ g CMV per 500 μ l of a lipid vesicle suspension.

In a preferred embodiment the lipid vesicles contain a mixture of the anionic phospholipid, DOPS, and a neutral lipid such as DOPE or DOPC. Other negatively charged phospholipids that can be used to make lipid vesicles include dioleoyl phosphatidic acid (DOPA) and dioleoyl phosphatidyl glycerol (DOPG). In a more preferred embodiment the neutral lipid is DOPC and the ratio of DOPS:DOPC is between 2:1 and 1:2 and is preferably about 1:1. The ratio of negatively charged to neutral lipid should be greater than 1:9 because the presence of less than 10% charged lipid results in instability of the lipid vesicles because of vesicle fusion.

A particular lipid vesicle formulation can be tested by using the formulation to transfect a target cell population with a plasmid of about 5.0 kb in length that expresses

some readily detectable product in the transfected target cell. Lipid vesicles can be used to transfect a cell with the plasmid if the plasmid is condensed with PEI at an imine:phosphate ratio of about 9-4:1. The capacity of the lipid vesicle formulation to transfect a cell with a plasmid is indicative of the formulation's capacity to introduce a CMV into a cell and effect a transmutation.

Certain lipids, particularly the polycationic lipids, can be toxic to certain cell lines and primary cell cultures. The formulation of the lipid vesicles should be adjusted to avoid such toxic lipids.

Ligands for clathrin-coated pit receptors can be introduced into the lipid vesicles by a variety of means. Cerebrosides, such as lactocerebroside or galactocerebroside can be introduced into the lipid mixture and are incorporated into the vesicle to produce a ligand for the asialoglycoprotein receptor.

In an alternative embodiment the lipid vesicle further comprises an integral membrane protein that inserts itself into the lipid bilayer of the vesicle. In a specific embodiment the protein is a fusogenic (F-protein) from the virus alternatively termed Sendai Virus or Hemagglutinating Virus of Japan (HVJ). The preparation and use of F-protein containing lipid vesicles to introduce DNA into liver, myocardial and endothelial cells have been reported. See, e.g., U.S. Patent No. 5,683,866, International Application PCT JP97/00612 (published as WO 97/31656). See also, Ramani, K., et al., 1996, FEBS Letters **404**, 164-168; Kaneda, Y., et al., 1989, J. Biol. Chem. **264**, 121126-12129; Kaneda, Y., et al., 1989, Science **243**, 375; Dzau, V.J., et al., Proc. Natl. Acad. Sci. **93**, 11421-11425; Aoki, M., et al., 1997, J.Mol.Cardiol. **29**, 949-959.

6.4 THE USE OF THE FORMULATIONS *IN VIVO*

The CMV of the invention can be parenterally administered directly to the target organ at a dose of between 50 and 250 $\mu\text{g/gm}$. When the target organ is the liver muscle or kidney, the CMV/macromolecular carrier complex can be injected directly into the organ. When the target organ is the liver, intravenous injection into the hepatic or portal veins of a liver, having temporarily obstructed circulation can be used. Alternatively the CMV/macromolecular complex can further comprise a hepatic targeting ligand, such as a

lactosyl or galactosyl saccharide, which allows for administration of the CMV/macromolecular complex intravenously into the general circulation.

When the target organ is the lung or a tissue thereof, e.g., the bronchiolar epithelium CMV/macromolecular complex can be administered by aerosol. Small particle aerosol delivery of liposomal/DNA complexes is described in Schwarz L.A., et al., 1996, Human Gene Therapy 7, 731-741.

When the target organ is the vascular endothelium, as for example in von Willebrand's Disease, the CMV/macromolecular complex can be delivered directly into the systemic circulation. Other organs can be targeted by use of liposomes that are provided with ligands that enable the liposome to be extravasated through the endothelial cells of the circulatory system.

For enzymatic defects, therapeutic effects can be obtained by correcting the genes of about 1% of the cells of the affected tissue. In a tissue in which the parenchymal cells have an extended life, such as the liver, treatments with CMV can be repeatedly performed to obtain an increased therapeutic effect.

7. EXAMPLES

7.1 CMV/MACROMOLECULAR CARRIER COMPLEXES

7.1.1 Lipid Nanospheres

Materials

Egg phosphatidylcholine (EPC), DOTAP and galactocerebroside (Gc) (Avanti Polar Lipids); soybean oil (Sigma Chemical Co.); dioctadecyldiamidoglycyl spermine (DOGS®) (Promega).

Methods

EPC, DOTAP and Gc were previously dissolved at defined concentrations in chloroform or anhydrous methanol and stored in small glass vials in desiccated containers at -20°C until use. EPC (40-45 mg), DOTAP (200 µg) and Gc (43 µg) solutions were aliquoted into a small 10 x 75 mm borosilicate tube and solvents removed under a stream of nitrogen. CMV were diluted in 0.15 M NaCl (~80-125 µg/250-300 µl); DOGS (as a 10 mg/ml solution in ethanol) was diluted into 250-300 µl 0.15 M NaCl at 3-5 times the weight of added CMV. The two solutions were mildly vortexed to mix contents and then CMV solution was added slowly to the DOGS solution. The contents were mixed by gentle tapping and inverting the tube a few times. The DOGS-complex solution was added to the dried lipids followed by soybean oil (40-45 mg), the mixture was vortexed on high for a few seconds and bath sonicated in a FS-15 (Fisher Scientific) bath sonicator for ~1 hr in a 4°C temperature controlled room. Occasionally, the tube was removed from the bath and vortexed. When a uniform looking, milky suspension was formed (with no obvious separation of oil droplets), it was extruded through a series of polycarbonate membranes down to a pore size of 50 nm. Preparations were stored at 4°C until use and vortexed before use.

7.1.2 Negatively Charged Targeted Lipid Vesicles

Materials

Dioleoyl phosphatidylcholine (DOPC), dioleoyl phosphatidylserine (DOPS), galactocerebroside (Gc) or lactosylcerebroside, (Avanti Polar Lipids).

Methods

DOPS, DOPC and Gc at a molar ratio of 1:1:0.16 (500 μg total lipid) were dissolved in chloroform:methanol (1:1 v/v) and then dried under a stream of nitrogen to obtain a uniform lipid film. The CMV were diluted in 500 μl of 0.15 M NaCl (approximately 100-250 $\mu\text{g}/500\mu\text{l}$). The solution was added to the lipid film at room temperature. Lipids were dispersed entirely by alternate mild vortexing and warming (in a water bath at 37-42°C). After a uniform milky suspension was formed, it was extruded through a series of polycarbonate membranes (pore sizes 0.8, 0.4, 0.2, 0.1 and 0.05 μm) using a Liposofast® mini-extruder. Extrusion was done 5 to 7 times through each pore size. After preparation, lipid vesicles were stored at 4°C until use. Under these conditions the lipid vesicles were stable for at least one month. The final product can be lyophilized.

7.1.3 Neutral Targeted Lipid Vesicles

Materials

Dioleoyl phosphatidylcholine (DOPC), dioleoyl phosphatidylethanolamine (DOPE), galactocerebroside (Gc) or lactosylcerebroside, (Avanti Polar Lipids).

Methods

DOPC, DOPE and Gc (1:1:0.16 molar ratio) or DOPC:Gc (1:0.08) were dissolved in chloroform:methanol (1:1 v/v) and then dried under a stream of nitrogen to obtain a uniform lipid film. The oligonucleotides (or chimeric molecules) were diluted in 500 μl of 0.15 M NaCl (approximately 100-250 $\mu\text{g}/500\mu\text{l}$). The solution was added to the lipid film at room temperature. Lipids were dispersed entirely by alternate mild vortexing and warming (in a water bath at 37-42°C). After a uniform milky suspension was formed, it was extruded through a series of polycarbonate membranes (pore sizes 0.8, 0.4, 0.2, 0.1 and 0.05 μm) using a Liposofast® mini-extruder. Extrusion was done 5 to 7 times through each pore size. After preparation, lipid vesicles were stored at 4°C until use. The size of the lipid vesicles of the preparation was stable for about 5 days.

7.1.4 Positively Charged Targeted Lipid Vesicles

Materials

Dioleoyl phosphatidylcholine (DOPC), dioleoyl trimethylammonium propane (DOTAP), galactocerebroside (Gc) or lactosylcerebroside, (Avanti Polar Lipids). Polyethylenimine (PEI) (M.W. 800 Kd), Fluka Chemicals.

Methods

DOPC, DOTAP and Gc (6:1:0.56 molar ratio) (500 μ g total lipid) were dissolved in chloroform:methanol (1:1 v/v) and then dried under a stream of nitrogen to obtain a uniform lipid film. PEI was diluted to a concentration of 45 mg/100 ml using water. pH of the solution was adjusted to ~7.6 using HCl. This PEI stock solution was prepared fresh each time and was equivalent to approximately 50 nmol amine/ μ l. CMV were diluted into 0.15 M NaCl at a concentration of ~125 μ g in 250 μ l. PEI was further diluted into 250 μ l 0.15 M NaCl so that approximately 4 moles of PEI amine were present per mole of oligonucleotide/chimeric phosphate. PEI solution was added dropwise to the CMV solution (both at room temperature) and vortexed for 5-10 minutes. The PEI-complex solution was then added to the lipid film and the lipids dispersed as described above. After a uniform milky suspension was formed, it was extruded through a series of polycarbonate membranes (pore sizes 0.8, 0.4, 0.2, 0.1 and 0.05 μ m) using a Liposofast® mini-extruder. Extrusion was done 5 to 7 times through each pore size. After preparation lipid vesicles were stored at 4°C until use. Under these conditions the lipid vesicles were stable for at least one month. For longer and improved stability the final product can be lyophilized.

7.1.5 Lactosylated-PEI/PEI Complexes

PEI (25 kDa) was purchased from Aldrich Chemical (Milwaukee, WI). PEI (800 kDa) was purchased from Fluka chemicals (Ronkonkoma, NY, USA). Lactosylation of the PEI was carried out by modification of a previously described method for the conjugation of oligosaccharides to proteins. Briefly, 3 to 5 ml of PEI (0.1 to 1.2 M_{monomer}) in ammonium acetate (0.2 M) or Tris buffer (0.2 M) (pH 7.6) solution was incubated with 7 to 8 mg of sodium cyanoborohydride (Sigma Chemical Co., St. Louis, MO) and

approximately 30 mg of lactose monohydrate (Sigma Chemical Co., St. Louis, MO). Reaction was carried out in polypropylene tubes, tightly capped in a 37°C shaking water bath. After 10 days the reaction mixture was dialyzed against distilled water (500 ml) for 48 h with 1 to 2 changes of water. The purified complex was sterile filtered through 0.2 μ m filter and stored at 4°C. The amount of sugar (as galactose) associated with PEI was determined by the phenol-sulphuric acid method.

The number of moles of free amine (primary + secondary) in the lactosylated PEI was determined as follows: a standard curve was set up using a 0.02M stock solution of PEI; several aliquots of the stock were diluted to 1ml using deionized water in glass tubes, then 50 μ l of Ninhydrin reagent (Sigma Chemical Co., St. Louis, Mo) was added to each tube and vortexed vigorously for 10 sec. Color development was allowed to proceed at room temperature for 10 to 12 min. and then O.D. was read (within 4 minutes) at 485 nm on a Beckman DU-64 spectrophotometer. 20 to 50 μ l aliquots of the L-PEI samples were treated as above and the number of moles of free amine was determined from the standard curve. Lactosylated-PEI (L-PEI) complexes were prepared as follows: an equivalent of 3 mmol of amine as L-PEI and 3 mmol of amine as PEI, per mmol of RNA/DNA phosphate, were mixed together and diluted in 0.15M NaCl as required; the mixture was added dropwise to a solution of the chimeric and vortexed for 5 min.

To verify complete association of the chimeric oligonucleotides with PEI or L-PEI, gel analysis (4% LMP agarose) of the uncomplexed and complexed chimerics was performed. To determine the degree of protection against nuclease degradation provided by complexation of the chimerics, samples were treated with RNase and DNase. After a chloroform phenol extraction, the complexes were dissociated using heparin (50 units/ μ g nucleic acid) and the products analyzed on a 4% LMP agarose gel.

7.2 DEMONSTRATION OF PEI/CMV MEDIATED ALTERATION OF RAT AND HUMAN FACTOR IX

Materials. Fetal bovine serum was obtained from Atlanta Biologicals, Inc. (Atlanta, GA). The terminal transferase, fluorescein-12-dUTP, Expand™ high fidelity PCR system, dNTPs and high pure PCR template preparation kit were obtained from Boehringer Mannheim Corp. (Indianapolis, IN). Reflection™ NEF-496 autoradiography

film and Reflection™ NEF-491 intensifying screens were from DuPont NEN® Research Products (Boston, MA). Polyethylenimine (PEI) 800 kDa was obtained from Fluka Chemical Corp. (Ronkonkoma, NY). The [γ -³²P]ATP was obtained from ICN Biochemicals, Inc. (Costa Mesa, CA). pCR™2.1 was obtained from Invitrogen (San Diego, CA). OPTIMEM™, Dulbecco's modified Eagle's medium, William's E medium and oligonucleotides 365-A and 365-C were from Life Technologies, Inc. (Gaithersburg, MD). Spin filters of 30,000 mol wt cutoff were purchased from Millipore Corp. (Bedford, MA). Dil and SlowFade™ antifade mounting medium were obtained from Molecular Probes, Inc. (Eugene, OR). T4 polynucleotide kinase was purchased from New England Biolabs, Inc. (Beverly, MA). MSI MagnaGraph membrane was purchased from Micron Separations, Inc. (Westboro, MA). The primers used for PCR amplification were obtained from Oligos Etc., Inc. (Wilsonville, OR). Tetramethylammonium chloride was purchased from Sigma Chemical Company (St. Louis, MO). All other chemicals were molecular biology or reagent grade and purchased from Aldrich Chemical Company (Milwaukee, WI), Curtin Matheson Scientific, Inc. (Eden Prairie, MN), and Fisher Scientific (Itasca, IL).

Oligonucleotide synthesis. Chimeric RNA/DNA oligonucleotides HIXF, RIXF and RIXR were synthesized. The CMV were prepared with DNA and 2'-O-methyl RNA phosphoramidite nucleoside monomers on an ABI 394 synthesizer. The DNA phosphoramidite exocyclic amine groups were protected with benzoyl (adenosine and cytidine) and isobutyryl (guanosine). The protective groups on the 2'-O-methyl RNA phosphoramidites were phenoxyacetyl for adenosine, isobutyryl for cytidine, and dimethylformamide for guanosine. The base protecting groups were removed following synthesis by heating in ethanol/concentrated ammonium hydroxide for 20 h at 55°C. The crude oligonucleotides were electrophoresed on 15% polyacrylamide gels containing 7 M urea, and the DNA visualized using UV shadowing. The chimeric molecules were eluted from the gel slices, concentrated by precipitation and desalted using G-25 spin columns. Greater than 95% of the purified oligonucleotides were full length.

The sequence of the wild type and "mutant" rat Factor IX are

(SEQ ID No. 27) 365
 wt AAA GAT TCA TGT GAA GGA GAT AGT GGG GGA CCC CAT GTT
 Lys Asp Ser Cys Glu Gly Asp Ser Gly Gly Pro His Val
 (SEQ ID No. 28)

(SEQ ID No. 29)
 mt AAA GAT TCA TGT GAA GGA GAT CGT GGG GGA CCC CAT GTT
 Arg

The structure of the RIXR, RIXF and HIXR CMV is as follows:

Chimeric Oligonucleotides

RIXR (SEQ ID No. 30)

TGCGCG-ccccagggggTGCTAgaggaaguguT

T T

T T

TCGCGC GGGGTCCCCCACGATCTCCTTCACAT

3' 5'

RIXR_c (SEQ ID No. 31)

TGCGCG-acacuuccucTAGCAccccccuggggT

T T

T T

TCGCGC TGTGAAGGAGATCGTGGGGGACCCCT

3' 5'

RIXF (SEQ ID No. 32)

TGCGCG-acacuuccucTAGCAccccccuggggT

T T

T T

TCGCGC TGTGAAGGAGATCGTGGGGGACCCCT

3' 5'

HIXF (SEQ ID No. 33)
 TGC GCG -acaguuccuc TAGCA ccccccuggggT
 T T
 T T
 TCGCGC TGTCAAGGAGAT TCGT GGGGGACCCCT
 3' 5'

Uppercase letters are deoxyribonucleotides, lower case letters are 2'OMe-ribonucleotides. The nucleotide of the heterologous region is underlined.

Cell Culture, transfections and hepatocyte isolation. HuH-7 cells were maintained in Dulbecco's modified Eagle's medium containing 10% (vol/vol) heat inactivated fetal bovine serum in a humidified CO₂ atmosphere at 37°C. Twenty four hours prior to transfection 1 x 10⁵ cells were plated per 35 mm culture dish. At the time of transfection, the cells were rinsed twice with OPTIMEM™ media and transfections were performed in 1 ml of the same media. Eighteen hours after transfection, 2 ml of Dulbecco's modified Eagle's medium containing 20% (vol/vol) heat inactivated fetal bovine serum was added to each 35 mm dish and the cells maintained for an additional 30 h prior to harvesting for DNA isolation. A PEI (800 kDa) 10 mM stock solution, pH 7.0, was prepared. Briefly, the chimeric oligonucleotides were transfected with 10 mM PEI at 9 equivalents of PEI nitrogen per chimeric phosphate in 100 µl of 0.15 M NaCl at final concentrations of either 150 nM (4 µg), 300 nM (8 µg) and 450 nM (12 µg). After 18 h, an additional 2 ml of medium was added and reduced the chimeric concentrations to 50 nM, 100 nM, and 150 nM, respectively, for the remaining 30 h of culture. HuH-7 vehicle control transfections utilized the same amount of PEI as was used in the HuIXF transfections, but substituted an equal volume of 10 mM Tris-HCl pH 7.6 for the oligonucleotides.

Primary rat hepatocytes were isolated from 250 g male Sprague-Dawley rats (Harlan Sprague-Dawley, Inc., Indianapolis, IN) by a two step collagenase perfusion as previously described (Fan et al., *Oncogene* 12:1909-1919, 1996, which is hereby incorporated by reference) and plated on Primaria™ plates at a density of 4 x 10⁵ cells per 35 mm dish. The cultures were maintained in William's E medium supplemented with 10% heat inactivated FBS, 26 mM sodium bicarbonate, 23 mM HEPES, 0.01 U/ml

insulin, 2 mM L-glutamine, 10 nM dexamethasone, 5.5 mM glucose, 100 U/ml penicillin and 100 U/ml streptomycin. Twenty four hours after plating, the hepatocytes were washed twice with the same medium and 1 ml of fresh medium added and the cells transfected using PEI/chimeric oligonucleotide complexes at the identical concentrations as for the HuH-7 cells. After 18 h, an additional 2 ml of the medium was added and the cells harvested 6 or 30 h later.

Direct injection of chimeric oligonucleotides into liver. Male Sprague-Dawley rats (~175 g) were maintained on a standard 12 h light-dark cycle and fed *ad libitum* standard laboratory chow. The rats were anesthetized, a midline incision made the liver exposed. A clamp was placed on the hepatic and portal veins as they enter the caudate lobe, and 75 μ g of the 1:9 chimeric/PEI complex was injected in a final volume of 250 - 300 μ l directly into the caudate lobe. The lobe remained ligated for 15 min and then blood flow was restored by removing the clamp. After suturing the incision the animals were allowed to recover from the anesthesia and given food and water *ad libitum*. Vehicle controls were done substituting an equal volume of Tris-HCl pH 7.6 for the chimeric oligonucleotides. Twenty-four and 48 h post-injection the animals were sacrificed, the caudate lobe removed and the tissue around the injection site dissected for DNA isolation. DNA was isolated and the terminal exon of the rat factor IX gene was amplified by PCR.

Nuclear uptake of the chimeric molecules. Chimeric duplexes were 3' end-labeled using terminal transferase and fluorescein-12-dUTP according to the manufacturer's recommendation, and were then mixed with unlabeled oligonucleotides at a 2:3 ratio. Transfections were performed as described above and after 24 h the cells were fixed in phosphate buffered saline, pH 7.4, containing 4% paraformaldehyde (wt/vol) for 10 min at room temperature. Following fixation, the cells were counterstained using a 5 μ M solution of Dil in 0.32 M sucrose for 10 min according to the manufacturer's recommendation. After washing with 0.32 M sucrose and then phosphate buffered saline, pH 7.4, the cells were coverslipped using SlowFade™ antifade mounting medium in phosphate buffered saline and examined using a MRC1000 confocal microscope (BioRad, Inc., Hercules, CA). The caudate lobes of liver *in situ* were injected with fluorescently-labeled chimerics as described above and harvested 24 h post-

injection. The lobes were bisected longitudinally, embedded using OCT and frozen. Cryosections were cut ~10 μ m thick, fixed for 10 min at room temperature using phosphate buffered saline, pH 7.4, containing 4% paraformaldehyde (wt/vol). Following fixation, the cells were counterstained using a 5 μ M solution of Dil in 0.32 M sucrose for 10 min according to the manufacturer's recommendation. After washing with 0.32 M sucrose and then phosphate buffered saline, pH 7.4, the sections were coverslipped using SlowFade™ antifade mounting medium and examined using a MRC1000 confocal microscope (BioRad, Inc.). The collection series for the fixed cells and sectioned tissue were made at 1 μ m steps to establish the presence of the chimeric in the nucleus.

DNA isolation and cloning. The cells were harvested by scrapping 24 and 48 h after transfection. Genomic DNA larger than 100-150 base pairs was isolated using the high pure PCR template preparation kit according to the manufacturer's recommendation. PCR amplification of a 317-nt fragment of the eighth exon in the human liver factor IX gene was performed with 500 ng of the isolated DNA. The primers used were 5'-CATTGCTGACAAGGAATACACGAAC-3' (SEQ ID No. 34) and 5'-ATTTCCTTTTCATTGCACACTCTTC-3' (SEQ ID No. 35) corresponding to nucleotides 1008-1032 and 1300-1324, respectively, of the human factor IX cDNA. Primers were annealed at 58°C for 20 sec, elongation was for 45 sec at 72°C and denaturation proceeded for 45 sec at 94°C. The sample was amplified for 30 cycles using Expand Hi-fidelity™ polymerase. PCR amplification of a 374-nt fragment of the rat factor IX gene was performed with 500 ng of the isolated DNA from either the primary hepatocytes or liver caudate lobe. The primers used were 5'-ATTGCCTTGCTGGAAGTGGATAAC-3' (SEQ ID No. 36) and 5'-TTGCCTTTTCATTGCACATTCTTCAC-3' (SEQ ID No. 37) corresponding to nucleotides 433-457 and 782-806, respectively, of the rat factor IX cDNA. Primers were annealed at 59°C for 20 sec, elongation was for 45 sec at 72°C and denaturation proceeded for 45 sec at 94°C. The sample was amplified for 30 cycles using Expand Hi-fidelity™ polymerase. The PCR amplification products from both the human and rat factor IX genes were subcloned into the TA cloning vector pCR™2.1 according to the manufacturer's recommendations, and the ligated material used to transform frozen competent *Escherichia coli*.

Colony hybridization and sequencing. Eighteen to 20 h after plating, the colonies were lifted onto MSI MagnaGraph nylon filters, replicated and processed for hybridization according to the manufacturer's recommendation. The filters were hybridized for 24 h with 17 mer oligonucleotide probes 365-A (5'-AAGGAGATAGTGGGGGA-3') (SEQ ID No. 38) or 365-C (5'-AAGGAGATCGTGGGGGA-3') (SEQ ID No. 39), where the underlined nucleotide is the target of the mutagenesis. The probes were ³²P-end-labeled using [γ -³²]ATP (> 7,000 Ci/mmol) and T4 polynucleotide kinase according to the manufacturer's recommendations. Hybridizations were performed at 37°C in 2X sodium chloride sodium citrate containing 1% SDS, 5X Denhardt's and 200 μ g/ml denatured sonicated fish sperm DNA. After hybridization, the filters were rinsed in 1X sodium chloride phosphate EDTA, 0.5% SDS and then washed at 54°C for 1 h in 50 mM Tris-HCl, pH 8.0 containing 3 M tetramethylammonium chloride, 2 mM EDTA, pH 8.0, 0.1% SDS. Autoradiography was performed with NEN® Reflection film at -70°C using an intensifying screen. Plasmid DNA was prepared from colonies identified as hybridizing with 365-A or 365-C using Qiagen minprep kit (Chatsworth, CA) and subjected to automatic sequencing using the mp13 reverse primer on an ABI 370A sequencer (Perkin-Elmer, Corp., Foster City, CA).

Results *In Vivo*

Chimeric oligonucleotides were fluorescein-labeled and used to determine whether direct injection into the caudate lobe of the liver was feasible. The results indicated that the hepatocytes adjacent to the injection site within the caudate lobe showed uptake of the fluorescently-labeled chimeric molecules similar to that observed in isolated primary hepatocytes and HuH-7 cells. Although some punctate material was present in the cytoplasm, the labeled material was detected primarily in the nucleus. In fact, only nuclear labeling was observed in hepatocytes farthest from the injection site. The unlabeled PEI/RIXF chimeric complexes and vehicle controls were injected directly into the caudate lobe using the same protocol and the animals sacrificed 24 and 48 h post-injection. Liver DNA was isolated as described in Methods, subjected to PCR amplification of a 374 nt sequence spanning the targeted nt exchange site. Following

subcloning and transformation of *Escherichia coli* with the PCR amplified material, duplicate filter lifts of the transformed colonies were performed. The filters were hybridized with ³²-labeled 17-mer oligonucleotides specific for either 365-A (wild-type) or 365-C (factor IX mutation) and processed post-hybridization as described in Methods. Rats which received direct hepatic injection of the RIXF chimeric molecules exhibited a A→C conversion frequency of ~1% at both 24 and 48 h. In contrast, the vehicle controls showed no hybridization with the 365-C probe. Colonies that hybridized with the 365-C probe from the RIXF treated animals were cultured, the plasmid DNA isolated and subjected to sequencing to confirm the A→C conversion. The ends of the amplified 374-nt fragment correspond exactly with the primers and the only nucleotide change observed was an A→C at the targeted exchange site.

7.3 DEMONSTRATION OF LACTOSYLATED-PEI/CMV MEDIATED ALTERATION OF RAT FACTOR IX

7.3.1 Results

CMV complexed to a mixture of lactosylated-PEI and PEI was prepared using the RIXR oligonucleotide as described in Section 6.1.5 above. A CMV directed to the complementary strand of the same region of the factor IX was also constructed (RIXR_C).

Conversion of the targeted nucleotide at Ser³⁶⁵ by the chimeric oligonucleotides

The nuclear localization of the fluorescently-labeled chimeric molecules indicated efficient transfection in the isolated rat hepatocytes. The cultured hepatocytes were then transfected with the unlabeled chimeric molecules factor RIXR_C and RIXR at comparable concentrations using 800 kDa PEI as the carrier. Additionally, vehicle control transfections were performed simultaneously. Forty eight hours after transfection, the cells were harvested and the DNA isolated and processed for hybridization as described in Section 6.1.5. The A→C targeted nucleotide conversion at Ser³⁶⁵ was determined by hybridization of duplicate colony lifts of the PCR-amplified and cloned 374-nt stretch of exon 8 of the factor IX gene (Sarkar, B., Koeberl, D. D. & Somer, S. S., "Direct Sequencing of the activation peptide and the catalytic domain of the factor IX gene in six species," *Genomics*, 6, 133-143, 1990.) The 17 mer oligonucleotide probes used to distinguish between the wild-type 365-A (5'-AAGGAGATAGTGGGGGA-3') (SEQ ID No.

40) or converted 365-C (5'-AAGGAGATCGTGGGGGA-3') (SEQ ID No. 41) corresponded to nucleotides 710 through 726 of the cDNA sequence.

The overall frequency of conversion of the targeted nucleotide was calculated by dividing the number of clones hybridizing with the 365-C oligonucleotide by the total number of clones hybridizing with both oligonucleotide probes. The results are summarized in Table III for RIXR_C. A→C conversion at Ser³⁶⁵ was observed only in primary hepatocytes transfected with the RIXR or RIXR_C. Similar conversion frequencies were observed in hepatocytes transfected with RIXR or RIXR_C. Neither vehicle transfected cells nor those transfected with other chimeric oligonucleotides yielded any clones hybridizing with the 365-C oligonucleotide probe (unpublished observations). Additionally, no hybridization of the 365-C oligonucleotide probe was observed to clones derived from DNA isolated from untreated hepatocytes and PCR-amplified in the presence of 0.5 to 1.5 µg of the oligonucleotides. The A→C conversion rate in the isolated hepatocytes was also dose dependent using lactosylated PEI derivatives as described in Section 6.1.5 and was as high as 19%. RT-PCR and hybridization analysis of RNA isolated from cultured cells transfected in parallel with lactosylated PEIs demonstrated A→C conversion frequencies ranging from 11.9 to 22.3%.

Site-directed nucleotide exchange by chimeric oligonucleotides in intact liver

The fluorescein-labeled oligonucleotides were also used to determine cellular uptake of the chimeric molecules after direct injection into the caudate lobe of the liver. The results indicated that hepatocytes adjacent to the injection site in the caudate lobe showed uptake of the fluorescently-labeled chimerics similar to that observed in the isolated rat hepatocytes. Although some punctate material was present in the cytoplasm of the hepatocytes, the labeled material was primarily present in the nucleus. In fact, only nuclear labeling was observed in those areas farthest from the injection site. The unlabeled RIXR chimeric oligonucleotides and vehicle controls were then administered *in vivo* by tail vein injection of the 25 kDa PEI and liver tissue harvested 5 days post-injection. Liver DNA was isolated and subjected to PCR amplification of a 374-nt sequence spanning the targeted nucleotide exchange site, using the same primers as those used with the primary hepatocytes. Following subcloning and transformation of *E. coli*

with the PCR-amplified material, duplicate filter lifts of the transformed colonies were done. The filters were hybridized with the same ^{32}P -labeled 17-mer oligonucleotides specific for either 365-A (wild-type) or 365-C (mutant) and processed post-hybridization. Rats treated with 100 μg of the RIXR chimeric oligonucleotides exhibited an A \rightarrow C conversion frequency ranging from 13.9% to 18.9%, while those that received a total of 350 μg in two injections showed 40% conversion. In contrast, the vehicle controls showed no hybridization with the 365-C probe. RT-PCR hybridization of isolated RNA indicated A \rightarrow C conversion frequencies of 26.4% to 28.4% in the high dose livers. The APTT for vehicle-treated rats ranged from 89.7% to 181.9% of control values ($131.84\% \pm 32.89\%$), while the APTT for the oligonucleotide-treated animals ranged from 48.9% to 61.7% ($53.8\% \pm 4.8\%$).

The APTT times for a 1/10 dilution of rat test plasma in Hepes buffer (50 mM Hepes/100 mM NaCl/0.02% NaN_3 , pH 7.4) were determined for both normal ($n = 9$) and the double injected animals ($n = 3$). The factor IX activity of duplicate samples was determined from a log-log standard curve that was constructed from the APTT results for dilution (1:10 to 1:80) of pooled plasma from 12 normal male rats, 6-8 weeks old. The APTT results for the normal rats ranged from 89.7% to 181.9% of the control values (mean = $131.84\% \pm 32.89\%$), while the APTT results for the double injected animals ranged from 49.0% to 61.7% (mean $53.8\% \pm 5.8\%$). The APTT clotting time in seconds for the normal rats ranged from 60.9 seconds to 81.6 seconds (mean = 71.3 ± 7.3 seconds) while the APTT times ranged from 92.3 seconds to 98.6 seconds (mean = 96.3 ± 2.9 seconds) for the double-infected rats.

Sequence analysis of the mutated factor IX gene in isolated hepatocytes and intact liver

Direct sequencing of the wild-type and mutated genes was performed to confirm the results from the filter hybridizations in both the *in vitro* and *in vivo* studies. At least 10 independent clones hybridizing to either 365-A or 365-C from the intact liver or isolated hepatocytes were analyzed. The results of the sequencing indicated that colonies hybridizing to 365-A (Fig. 6, top panel) exhibited the wild-type IX sequence, i.e. an A at Ser³⁶⁵ of the reported cDNA sequence. In contrast, those colonies derived from the factor RIXR_C transfected primary hepatocytes hybridizing to the 365-C oligonucleotide probe

converted to a C at Ser³⁶⁵. The same A→C conversion at Ser³⁶⁵ was observed in the clones derived from the transfected rat liver that hybridized with the 17 mer 365-C oligonucleotide probe. The entire 374-nt PCR amplified region of the factor IX gene was sequenced for all the clones and no alteration other than the indicated changes at Ser³⁶⁵ was detected. Finally, the start and end points of the 374-nt PCR amplified genomic DNA derived from both the primary hepatocytes and the intact liver corresponded exactly to those of the primers used for the amplification process, indicating that the cloned and sequenced DNA was derived from genomic DNA rather than nondegraded chimeric oligonucleotides.

Table III Percent A-C conversion at Ser³⁶⁵ of rat factor IX genomic DNA by colony lift hybridizations

PEI Deliver System		365-C clones	Total clones	A-C (%)
PEI 800 kDa ¹	<u>Concentration</u>			
<i>In vitro</i>	150 nM	24	572	4.2
	300	31	367	8.5
	450	63	502	12.5
Lac-PEI 800 kDa				
<i>In vitro</i>	90	18	337	5.3
	180	34	300	11.3
	270	47	253	18.6
Lac-PEI 25 kDa				
<i>In vitro</i>	90	28	527	5.3
	180	53	417	12.7
	270	60	305	19.7
Lac-PEI 25 kDa ²	<u>Dose</u>			
<i>In vivo</i> x1	100 µg	24	166	14.5
		71	386	18.4
		50	360	13.9
Lac-PEI 25 kDa				
<i>In vivo</i> x2	350 µg	237	601	39.4
		228	563	40.5
		271	678	40.0

¹The data shown for the primary hepatocyte transfections represents a mean of two experiments.

²The *in vivo* chimeric/PEI complexes were administered in a volume of 300 µl of 5% dextrose by tail vein injection. The results of three animals at each dose are shown individually.

7.3.2 Materials and Methods

In vivo delivery of the chimeric oligonucleotides. Male Sprague-Dawley rats (Harlan Sprague-Dawley, Inc.) (~50 g) were maintained on a standard 12 h light-dark cycle and fed *ad libitum* standard laboratory chow. Vehicle controls and lactosylated 25 kDa PEI at a ratio of 6 equivalents of PEI nitrogen per chimeric phosphate were administered in 300 μ l of 5% dextrose (Abdallah, B. et al., "A powerful nonviral vector for *in vivo* gene transfer into the adult mammalian brain: polyethylenimine; *Human Gene Therapy*, 7, 1947-1954, 1996.). The aliquots were administered by tail vein injection either as a single dose of 100 μ g or divided dose of 150 μ g and 200 μ g on consecutive days. Five days post-injection, liver tissue was removed for DNA and RNA isolation. DNA was isolated as previously described (Kren, B. T., Trembley, J. H. & Steer, C. J., "Alterations in mRNA stability during rat liver regeneration," *Am. J. Physiol.*, 270, G763-G777, 1996) for PCR amplification of exon 8 of the rat factor IX gene. RNA was isolated for RT-PCR amplification of the same region as the genomic DNA using RNAexol and RNAmate (Intermountain Scientific Corp., Kaysville, UT) according to the manufacturer's protocol.

Factor IX activity assay. Blood samples from vehicle (n = 9) and oligonucleotide-treated (n = 3) rats were collected 20 days after the second tail vein injection in 0.1 vol. of 0.105 M sodium citrate/citric acid. After centrifugation at 2,500 x g and then 15,000 x g the resulting plasma was stored at -70°C. The factor IX activity was determined from activated partial thromboplastin time (APTT) assays. Briefly, 50 μ l of APTT reagent (DADE, Miami, FL), 50 μ l of human factor IX-deficient plasma (George King Biomedical, Overland, KS), and 50 μ l of 1/10 dilution of rat test plasma in Hepes buffer (50 mM Hepes/100 mM NaCl/0.02% NaN₃, pH 7.4) were incubated at 37°C for 3 min in an ST4 coagulometer (American Bioproducts, Parsippany, NJ). Clotting was initiated by addition of 50 μ l of 33 mM CaCl₂ in Hepes buffer. Factor IX activity of duplicate samples was determined from a log-log standard curve constructed from the APTT results for dilution (1:10 to 1:80) of pooled plasma from normal male rats (n = 12).

DNA/RNA isolation and cloning. The cells were harvested by scrapping 48 h after transfection. Genomic DNA larger than 100-150 base pairs was isolated using the high

pure PCR template preparation kit (Boehringer Mannheim, Corp., Indianapolis, IN). RNA was isolated using RNeasyTM B (Tel-Test, Inc., Friendswood, TX), according to the manufacturer's protocol. PCR amplification of a 374-nt fragment of the rat factor IX gene was performed with 300 ng of the isolated DNA from either the primary hepatocytes or liver tissue. The primers were designed as 5'-ATTGCCTTGCTGGAAGTGGATAAAC-3' (SEQ ID No. 42) and 5'TTGCCTTTTCATTGCACATTCTTCAC-3' (SEQ ID No. 43) (Oligos Etc., Wilsonville, OR) corresponding to nucleotides 433-457 and 782-806, respectively, of the rat factor IX cDNA. Primers were annealed at 59°C for 20 sec, elongation was for 45 sec at 72°C and denaturation proceeded for 45 sec at 94°C. The sample was amplified for 30 cycles using Expand Hi-fidelityTM polymerase (Boehringer Mannheim, Corp.). The PCR amplification products from both the hepatocytes and intact liver factor IX genes were subcloned into the TA cloning vector pCRTM2.1 (Invitrogen, San Diego, CA), and the ligated material used to transform frozen competent *E. coli*. To rule out PCR artifacts 300 ng of control DNA was incubated with 0.5, 1.0 and 1.5 µg of the oligonucleotide prior to the PCR-amplification reaction. Additionally, 1.0 µg of the chimeric alone was used as the "template" for the PCR amplification.

RT-PCR amplification was done utilizing the TitanTM one tube RT-PCR system (Boehringer Mannheim, Corp.) According to the manufacturer's protocol using the same primers as those used for the DNA PCR amplification. To rule out DNA contamination, the RNA samples were treated with RQ1 DNase free RNase (Promega Corp., Madison, WI) and RT-PCR negative controls of RNased RNA samples were performed in parallel with the RT-PCR reaction. Each of the PCR reactions were ligated into the same TA cloning vector and transformed into frozen competent *E. coli*.

Colony hybridization and sequencing. Eighteen to 20 h after plating, the colonies were lifted onto MSI MagnaGraph nylon filters, replicated and processed for hybridization according to the manufacturer's recommendation. The filters were hybridized for 24 h with 17 mer oligonucleotide probes 365-A (5'AAGGAGATAGTGGGGGA-3') (SEQ ID No. 44) OR 365-C (5'-AAGGAGATCGTGGGGGA-3') (SEQ ID No. 45) (Life technologies, Inc., Gaithersburg, MD), where the underlined nucleotide is the target for

mutagenesis. The probes were ^{32}P -end-labeled using (γ - ^{32}P) ATP ($> 7,000$ Ci/mmol) and T4 polynucleotide kinase (New England Biolabs, Inc., Beverly MA). Hybridizations were performed at 37°C in 2X sodium chloride sodium citrate containing 1% SDS, 5X Denhardt's and $200\text{ }\mu\text{g/ml}$ denatured sonicated fish sperm DNA. After hybridization, the filters were rinsed in 1X sodium chloride sodium phosphate EDTA, 0.5% SDS and then washed at 54°C for 1 h in 50 mM Tris-HCl, pH 8.0 containing 3 M tetramethylammonium chloride, 2 mM EDTA, pH 8.0, 0.1% SDS (Melchior, W. B. & Von Hippel, P. H. "Alteration of the relative stability of dA.dT and dG.dC base pairs in DNA," Proc. Natl. Acad. Sci. USA, **70**, 298-302, 1973.). Autoradiography was performed with NEN[®]Reflection film at -70°C using an intensifying screen. Plasmid DNA was prepared from colonies identified as hybridizing with 365-A or 365-C using Qiagen miniprep kit (Chatsworth, CA) and subjected to automatic sequencing using the mp13 forward and reverse primers as well as a gene specific primer, 5'GTTGACCGAGCCACATGCCTTAG-3' (SEQ ID No. 46) corresponding to nucleotides 616 to 638 of the rat factor IX cDNA using an ABI 370A sequencer (Perkin-Elmer, Corp., Foster City, CA).

7.4 EXAMPLES OF CMV USEFUL FOR THE REDUCTION OF LDL LEVELS IN HUMANS

A CMV suitable for the modification of Apo B having a sequence comprising the sequence of SEQ ID No: 5 is given below.

Apo B 41/UR (mut→WT)

(SEQ ID No. 47)

```

u GCGCG gac ccg acc gaa uuc ggu aac ugu au
u                                     u
u                                     u
u CGCGC CTG GGC TGG CTT AAG CCA TTG ACA Tu
      3' 5'

```

A CMV suitable for the modification of Apo B having a sequence comprising the sequence of SEQ ID No: 12 is given below.

Apo B 5/U88 (mut→WT)

(SEQ ID No. 48)

```

u GCGCG cug uuc aaa gug uaC GGA TCC ucu uug acu gac gau
u
u
u CGCGC GAC AAG TTT CAC ATG CCT AGG AGA AAC TGA CTG CTu
      3' 5'

```

7.5 CORRECTION OF A CRIGLER-NAJJAR-LIKE MUTATION IN THE GUNN RAT

Mutant rats with hyperbilirubinemia, termed Gunn rats, have a single nucleotide deletion in the gene encoding bilirubin-uridinediphosphoglucuronate glucuronosyltransferase (*UGT1A1*). Roy Chowdhury, J., et al., 1991, J. Biol. Chem. **266**, 18294. Human patients with Crigler-Najjar syndrome type I also have mutations of the *UGT1A1* gene, resulting in life-long hyperbilirubinemia and consequent brain damage. Bosma, P.J., et al., 1992, FASEB J. **6**, 2859; Jansen, P.L.M., et al., Progress In Liver Diseases, **XIII**, Boyer, J.L., & Ockner, R.K., editors (W.B. Saunders, Phil. 1995), pp 125-150. The structure of CN3, a CMV designed to correct the Gunn rat mutation is given below.

CN3 (mut→WT)

(SEQ ID No. 49)

```

T GCGCG gg gac uua caG GAC CTT TAC uga ctt cua T
T
T
T CGCGC CC CTG AAT GTC CTG GAA ATG ACT GCC GAT T
      3' 5'

```

Gunn rat primary cultured hepatocytes were treated with 150 nM CN3 according to the above protocol except that the carrier was either the negatively charged glycosylated lipid vesicles of section 6.2.2 or a lactosylated-PEI carrier at a ratio of

oligonucleotide phosphate to imine of 1:4 . The results were 8.5% conversion with the negatively charged liposome and 3.6% conversion with lactosylated-PEI carrier.

Gunn rats were injected with 1 mg/Kg of CN3 complexed with either 25 kDa Lac-PEI or complexed with negatively charged Gc lipid vesicles (Gc-NLV) as described above. The rate of gene conversion was determined by cloning and hybridization according to the procedure described for factor IX. The results shown below indicate that between about 15% and 25% of the copies of the *UGT1A1* gene were converted.

**Frequency of Insertion of G at nucleotide 1239 of the UGT-1 Gene
(In Gunn Rats)**

Vehicle	Dosage	G Clones/Total Clones	Frequency (%)
Gc-NLV	1 mg	112/815	15.4
		208/761	27.3
		185/974	18.9
		39/273	14.6 ¹
		78/403	19.3 ²
25 kDa PEI (Lactosylated)	1 mg	188/838	22.4
		254/1150	22.1
		245/997	24.6

¹Initial conversion frequency determined.

²Conversion frequency determined 7 days after 70% partial hepatectomy.

A Gunn rat was injected on five successive days with 1 mg/Kg of CN3 complexed with 25 kDa Lac-PEI as above. Twenty five days after the final injection the serum bilirubin had declined from 6.2 mg/dl to 3.5 mg/dl and remained at that level for a further 25 days.

7.6 CORRECTION OF A FACTOR IX MUTATION IN DOG

The Chapel Hill strain of dogs, which has a (G→A)¹⁴⁷⁷ mutation that results in hemophilia in the animals, was used to obtain primary cultured hepatocytes. Four CMV to correct this mutation have been synthesized.

DIX1 (mut→WT) (SEQ ID No. 50)

```

T gcgcg auu caa aga aTT GAC CCT AAT AAT cga ccc cT
T
T
T CGCGC TAA GTT TCT TAA CTG GGA TTA TTA GCT GGG GT
      3' 5'

```

DIX2 (mut→WT) (SEQ ID No. 51)

```

T gcgcg caa aga auu gAC CCT AAT aaU cga cT
T
T
T CGCGC GTT TCT TAA CTG GGA TTA TTA GCT GT
      3' 5'

```

DIX3 (mut→WT) (SEQ ID No. 52)
 u gcgcg auu caa aga auu gac ccu aau aau cga ccc cu
 u
 u
 u CGCGC TAA GTT TCT TAA CTG GGA TTA TTA GCT GGG Gu
 3' 5'

DIX4 (mut→WT) (SEQ ID No. 53)
 u gcgcg auu caa aga auu gac uccu aau aau cga ccc cu
 u
 u
 u CGCGC TAA GTT TCT TAA CTG GGA TTA TTA GCT GGG Gu
 3' 5'

DIX1 differs from DIX3 by the replacement of the intervening DNA segment with 2'-O-methyl RNA and replacement of the tetrathymidine linkers with tetrauracil. DIX 4 differs from DIX3 in that the mutational vector contains a mismatch in the mutator region. In DIX4 the 5' (lower) strand encodes the desired (wild-type) sequence while the 3' (upper) strand has the sequence of the target, i.e., the mutant sequence.

The hepatocytes were treated with 360 nM DIX1 complexed in either 25 kDa Lac-PEI or galactocerebroside-containing aqueous-cored, negatively charged lipid vesicles (Gc-NLV). The results are given in the table below.

Frequency of conversion of A to G at nucleotide 1477 of the Factor IX Gene
(Primary Hepatocytes from the Chapel Hill Strain of Hemophilia B Dogs)

Vehicle	Number of Times Transfected	Concentration	G Clones/Total Clones	Frequency (%)
Gc-NLV	Once	360 nM	30/195	15.44
			30/218	13.76
	Twice		30/118	25.4
Lac-PEI 25 kDa	Once*	360 nM	20/141	14.2
			48/348	13.3
	Twice		21/107	19.6

*RT-PCR on parallel transfected cultures gave an A to G conversion frequency of 11.1%

Each of the DIX2-DIX4 were also tested on primary cultured dog hepatocytes as above. The results showed that DIX2 worked poorly, possibly due to the low (25%) GC percentage. The subsequent experiments the results of DIX3 were about 16% conversion, while a parallel experiments DIX1 gave 10% conversion and the results of DIX4 were about as good as DIX1.

GenBank Sequence References for the Exons of the Human Apolipoprotein B-100 Gene

TABLE II

Exon No.	cDNA Boundary	GenBank Accession No. Sequence Reference
1	126 to 207	M19808
2	208 to 246	M19808
3	247 to 362	M19809
4	363 to 508	M19810
5	509 to 662	M19811
6	663 to 818	M19812
7	819 to 943	M19813
8	944 to 1029	M19813
9	1030 to 1249	M19815
10	1250 to 1477	M19816
11	1478 to 1595	M19818
12	1596 to 1742	M19818
13	1743 to 1954	M19820
14	1955 to 2192	M19820
15	2193 to 2359	M19821
16	2360 to 2561	M19823
17	2562 to 2729	M19824
18	2730 to 2941	M19824
19	2942 to 3124	M19825
20	3125 to 3246	M19825
21	3247 to 3457	M19827
22	3458 to 3633	M19828
23	3634 to 3821	M19828
24	3822 to 3967	M19828
25	3968 to 4341	M19828
26	4342 to 11913	M19828
27	11914 to 12028	M19828
28	12029 to 12212	M19828
29	12213 to 13816	M19828

SEQ ID	TABLE I	G/C#	NA Change	AA Change	AA	%APOB100	Restriction Site
No.	Sequence (5'→3')						
4	AGTCTGGATGGGIAAGCCGCCCTCA	15	A→T	K→Stop	1701	36.9	None
5	CTGGGCTGGCTTAAGCCATTGACAT	13	C→A	S→TAA	1876	40.8	+CTTAAG
6	GCTCTCTGGGGAIAACATACTGGGC	14	G→T	E→Stop	1921	41.8	None
7	GATGCCGTTGAGIAGCCCCCAAGAAT	13	A→T	K→Stop	2047	44.5	None
8	GAGAGGAATCGAIAAACCATTTATAG	10	C→T	Q→Stop	2085	45.4	+ATCGAT
9	TGTAAGAAAAATAAGAGAGAGCCCTG	10	C→A	Y→Stop	2110	45.9	None
10	GCAGCCCTGGGAIAACTCCACAGC	16	A→T	K→Stop	2116	46.0	None
11	GCAAGCTAATGATTAGCTGAAATTCATCAAT	8	T→G	Y→Stop	2124	46.2	+AGCT
12	CAAGTTTCACATGCCIAAGGAGAAACTGACTG	11	A→T	K→Stop	2138	46.5	+CCTAGG
13	ATATACAAATTCGATGAGATGATGCCAAAAT	9	T→G	L→Stop	2159	47.0	+CATG
14	AAACTATCTCAACTGIAAGACATATATGATAC	8	C→T	Q→Stop	2174	47.3	-CTGCAG
15	GCTAATATTATTGATIAAAATCATTGAAATTA	3	G→T	E→Stop	2204	48.0	+TTAA
16	TGATGAGCACTAGCAGATATCCGTGTA	11	T→G	Y→Stop	2216	48.3	+CTAG
17	CTGCAGCAGCTTIAAGAGACACATAC	12	A→T	K→Stop	2270	49.4	-CTTAAG
18	AACAGTGAGCTGIAAGTGGCCCGTTC	14	C→T	Q→Stop	2684	58.6	None
19	CAGACTTCCGTTAACCAGAAATCGC	12	T→A	L→Stop	2712	59.2	+GTTAAC
20	AAAGGTCATGGIAATGGGCTGCC	14	A→T	K→Stop	2930	64.0	None
21	ACATATATGATAIAAATTTGATCAGT	5	C→T	Q→Stop	2180	47.5	Physiologic

Table III

Seq ID No.	Sequence (5'→3')	G/C#	AA Change	NA Change	AA	Gene
22	ATGGAGGACGTG I GGGGCCGCTGG	18	R→C	C→T	112	Apo E
23	GACCTGCAGAAAG I GCCCTGGCAGTGT	15	R→C	C→T	158	Apo E
24	GACCTGCAGAAAG C CCCTGGCAGTGT	16	C→R	T→C	158	Apo E
25	TAAGGTCAGGAG I TTGAGACCAGCC	13	NA	A→T	491	Apo E
54	GGCGAGGACATG I GCGACCGGGCGC	19	R→C	C→T	149	Apo AI
55	GAGATGCGCGAC I GCGCGCGCGCCC	20	R→C	C→T	151	Apo AI
56	CGCGACCGCG C IGCGCGCATGTGG	20	R→C	C→T	153	Apo AI
57	AGCGACCAGCTG I GCCAGCGCTTGG	17	R→C	C→T	171	Apo AI
58	GAGCTGCGCCAG I GCTTGGCCCGCGC	19	R→C	C→T	173	Apo AI

WE CLAIM

1. A method of reducing LDL in the blood of a subject comprising altering an Apo B gene of a hepatocyte of the subject such that the transcript of the altered Apo B gene contains an in-frame stop codon whereby the altered gene encodes a protein having at least 1433 amino acids and not more than 3974 amino acids.
2. The method of claim 1, which further comprises the steps of determining the effect on the level of LDL of the alteration of the Apo B genes in the subject and subsequently adjusting the number of altered Apo B genes in the subject.
3. The method of claim 1, wherein the altered gene encodes a protein having at least 1841 amino acids and not more than 2975 amino acids.
4. The method of any one of claims 1-3, wherein the altered gene encodes a protein having a sequence of a fragment of SEQ ID No. 1, which fragment is at least amino acids 1-1841 and not more than amino acids 1-2975.
5. The method of any one of claims 1-4 which comprises administering a recombinagenic oligonucleobase which comprises a first and a second homologous region each having a sequence of at least 10 nucleobases selected from nt 4342-11913 of SEQ ID No: 2, whereby the alteration of the Apo B gene is effected.
6. The method of any one of claims 1-5, wherein the subject's fasting LDL serum cholesterol is reduced to below 140 mg/dl.
7. A composition for the modification of a human Apo B gene comprising an oligonucleobase which oligonucleobase comprises:
 - a. a first and a second homologous region that are each at least 8 nucleobases in length and together at least 20 nucleobases in length, which homologous regions are each homologous with a fragment of the sequence of nt 5649-9051 of SEQ ID No. 2, and

- b. a heterologous region that is disposed between the first and second homologous region,
such that the introduction of the sequence of the heterologous region into the Apo B gene results in the truncation of the protein encoded thereby.
- 8. The composition of claim 7, in which the first and the second homologous regions each comprises at least 3 contiguous nucleobase-pairs of hybrid-duplex.
- 9. The composition of claim 7 or 8, in which the sum of the lengths of the first and second homologous regions is not more than 60 nucleobases in length.
- 10. The composition of any one of claims 7-9, in which the homologous regions together comprise between 9 and 25 nucleobase pairs of hybrid-duplex.
- 11. The composition of any one of claims 7-10, in which the GC fraction of each homologous region is at least 33%.
- 12. The composition of any one of claims 7-10, in which the GC fraction of a homologous region is at least 50%.
- 13. The composition of any one of claims 7-10, in which the sequence of the oligonucleobase comprises the sequence of at least a 21 nucleobase fragment of any one of SEQ ID No. 4-21 or the complement thereof.
- 14. The composition of any one of claims 7-10, in which the sequence of the oligonucleobase comprises the sequence of at least a 25 nucleobase fragment of any one of SEQ ID No. 4-21 or the complement thereof.
- 15. The composition of any one of claims 7-14 which further comprises:
 - a. an aqueous carrier; and
 - b. a macromolecular carrier selected from the group consisting of

- i. an aqueous-cored lipid vesicle, wherein the aqueous core contains the oligonucleobase,
 - ii. a lipid nanosphere, which comprises a lipophilic salt of the oligonucleobase, and
 - iii. a polycation having an average molecular weight of between 500 daltons and 1.3 Md wherein the polycation forms a salt with the oligonucleobase.
16. The composition of claim 15, in which the macromolecular carrier further comprises a ligand for a clathrin-coated pit receptor.
17. The composition of claim 16, in which the clathrin-coated pit receptor is selected from the group consisting of the receptors for transferrin, nicotinic acid, carnitine, insulin and insulin like growth factor-1.
18. The composition of claim 16, in which the clathrin-coated pit receptor is an asialoglycoprotein receptor.
19. The composition of any one of claims 15-18, in which the macromolecular carrier is a negatively charged, aqueous-cored lipid vesicle.
20. The composition of any one of claims 16-18, in which the aqueous-cored lipid vesicle further comprises a cerebroside.
21. The composition of any one of claims 15-20, in which the aqueous-cored lipid vesicle comprises dioleoylphosphatidylcholine and dioleoylphosphatidylserine.
22. The composition of any one of claims 16, 18, 19 and 21 wherein the macromolecular carrier is an aqueous-cored lipid vesicle that comprises a fusogenic F-protein.

23. The composition of any one of claims 15-22, in which the oligonucleobase comprises:
- a. a first and a second homologous region that are together at least 16 and not more than 60 nucleobases in length, which regions are homologous with a target gene of a mammal; and
 - b. a heterologous region that is disposed between the first and second homologous region and is at least 1 and not more than 20 nucleobases in length, which is heterologous with the target gene and which contains the alteration.
24. A method of treatment and/or prophylaxis in a subject comprising altering an Apo E gene of a hepatocyte of the subject by introducing a substitution selected from the group (Arg→Cys)¹¹², (Arg→Cys)¹⁵⁸ and (Cys→Arg)¹⁵⁸.
25. The method of claim 24, wherein the subject is homozygous for Apo E4 and the alteration comprises the substitution (Arg→Cys)¹¹².
26. The method of claim 24 or 25, which comprises administering a chimeric mutational vector having a sequence which comprises SEQ ID No: 22.
27. The method of claim 24, wherein the treatment or prophylaxis comprises reducing the subject's fasting serum LDL cholesterol level and the alteration comprises the substitution (Arg→Cys)¹⁵⁸.
28. The method of claim 24 or 27, which comprises administering a chimeric mutational vector having a sequence which comprises SEQ ID No: 23.
29. The method of claim 24, wherein the subject suffers from Type III hyperlipidemia and the alteration comprises the substitution (Cys→Arg)¹⁵⁸.

30. The method of claim 24 or 29, which comprises administering a recombinagenic oligonucleobase having a sequence which comprises SEQ ID No: 24.
31. A composition for the alteration of a human Apo E gene comprising a recombinagenic oligonucleobase having a sequence comprising the sequence of at least a 21 nucleobase fragment of any one of SEQ ID No. 22 - 25 or the complement thereof.
32. A method of ameliorating atherosclerosis in a subject comprising altering an Apo A1 gene of a hepatocyte of the subject such that the altered Apo A1 protein forms dimers.
33. The method of claim 32, which further comprises the steps of determining the effect on the level of HDL of the alteration of the Apo A1 genes in the subject and subsequently adjusting the number of altered Apo A1 genes in the subject.
34. The method of claim 32, wherein the altered gene encodes a protein having a cysteine for arginine substitution at a position selected from the group consisting of residue 149, 151, 153, 171 and 173.
35. The method of claim 34, wherein the method comprises the administration of a recombinagenic oligonucleobase having a sequence comprising the sequence of at least 20 nucleotides of SEQ ID No. 54, No. 55, No. 56, No. 57 and No. 58.
36. The method of claim 35, which comprises the step of administering to the subject a composition comprising:
 - a) the recombinagenic oligonucleobase;
 - b) an aqueous carrier; and
 - c) a macromolecular carrier selected from the group consisting of

- (i) an aqueous-cored lipid vesicle, wherein the aqueous core contains the oligonucleobase,
 - (ii) a lipid nanosphere, which comprises a lipophilic salt of the oligonucleobase, and
 - (iii) a polycation having an average molecular weight of between 500 daltons and 1.3 Md wherein the polycation forms a salt with the oligonucleobase.
- 37. The method of claim 36, wherein the macromolecular carrier is a negatively charged, aqueous-cored lipid vesicle.
- 38. The method of claim 36, wherein the macromolecular carrier further comprises a ligand for a clathrin-coated pit receptor.
- 39. The method of claim 38, wherein the clathrin-coated pit receptor is an asialoglycoprotein receptor.
- 40. The method of claim 32, which comprises administering to the subject a composition comprising:
 - a) a recombinagenic oligonucleobase;
 - b) an aqueous carrier; and
 - c) a macromolecular carrier selected from the group consisting of
 - (i) an aqueous-cored lipid vesicle, wherein the aqueous core contains the oligonucleobase,
 - (ii) a lipid nanosphere, which comprises a lipophilic salt of the oligonucleobase, and
 - (iii) a polycation having an average molecular weight of between 500 daltons and 1.3 Md wherein the polycation forms a salt with the oligonucleobase.
- 41. The method of claim 40, wherein the macromolecular carrier is a negatively charged, aqueous-cored lipid vesicle.

42. The method of claim 40, wherein the macromolecular carrier further comprises a ligand for a clathrin-coated pit receptor.
43. The method of claim 42, wherein the clathrin-coated pit receptor is an asialoglycoprotein receptor.
44. A composition for the modification of a human Apo A1 gene comprising an oligonucleobase which oligonucleobase contains a sequence which comprises a fragment having the sequence selected from the group consisting of SEQ ID No. 54, No. 55, No. 56, No.57 and No. 58.
45. The composition of claim 44, which further comprises:
 - a) an aqueous carrier; and
 - b) a macromolecular carrier selected from the group consisting of
 - (i) an aqueous-cored lipid vesicle, wherein the aqueous core contains the oligonucleobase,
 - (ii) a lipid nanosphere, which comprises a lipophilic salt of the oligonucleobase, and
 - (iii) a polycation having an average molecular weight of between 500 daltons and 1.3 Md wherein the polycation forms a salt with the oligonucleobase.
46. The composition of claim 45, in which the macromolecular carrier further comprises a ligand for a clathrin-coated pit receptor.
47. The composition of claim 46, in which the clathrin-coated pit receptor is selected from the group consisting of the receptors for transferrin, nicotinic acid, carnitine, insulin and insulin like growth factor-1.
48. The composition of claim 46, in which the clathrin-coated pit receptor is an asialoglycoprotein receptor.

49. The composition of any one of claims 45-48, in which the macromolecular carrier is a negatively charged, aqueous-cored lipid vesicle.
50. The composition of any one of claims 46-48, in which the aqueous-cored lipid vesicle further comprises a cerebroside.
51. The composition of any one of claims 45-50, in which the aqueous-cored lipid vesicle comprises dioleoylphosphatidylcholine and dioleoylphosphatidylserine.
52. The composition of any one of claims 46, 48, 49 and 51 wherein the macromolecular carrier is an aqueous-cored lipid vesicle that comprises a fusigenic F-protein.

1/11





FIG.1

2/11

FIG. 2A

1 cctatccctgggggagggggcgggacaggggagccctataattggacaagtctgggatcccttgagtcctACTCAGCCCCAG
83 CGGAGGTGAAGGACGTCCTTCCCCAGGAGCCGgtgagaagcgagctcgggggcacggggatgagctcaggggcctctagaaa
165 gagctgggaccctgggaagccctggcctccaggtagtctcaggagagctactcggggtcgggcttggggagaggaggagcgg
247 gggtgaggcaagcagcaggggactggacctgggaagggctgggcagcagagacgacccgacccgctagaagggtgggtgggg
329 agagcagctggactgggatgtaagccatagcaggactccacgagttgtcactatcattatcgagcacctactgggtgtcccc
411 agtgtcctcagatctccataactggggagccaggggcagcgacacggtagctagccgtcgattggagaactttaaatgagg
493 actgaattagctcataaatggaacacggcgcttaactgtgaggttggagcttagaatgtgaagggagaatgaggaatgcgag
575 actgggactgagatggaacccggcggtggggaggggtgggggatggaatttgaacccgggagaggaagatggaattttct
657 atggaggccgacctggggatggggagataagagaagaccaggagggttaaatagggaatgggttggggcggttggtaa
739 atgtgctgggattaggctgttgcagataatgcaacaaggcttggaaaggctaacctggggtgaggccgggttggggcgctgg
821 ggggtgggaggagtcctcactggcggttgattgacagtttctccttccccagACTGGCCAATCACAGGCAGGAAGATGAAGGT
903 TCTGTGGGCTGCGTTGCTGCTCACATTCTGGCAGgtatgggggcggggcttgctcggttccccccgctcctccccctctca
L W A A L L V T F L A M K V
985 tcctcacctcaacctcctggccccattcagacagacctggggccccctcttctgaggcttctgtgctgcttctggtctga
1067 acagcgatttgacgctctctgggcctcggtttccccatccttgagataggagttagaagttgttttgttgttgttgtt
1149 tgttgttgttttgttttttggatgaagtctcgctctgtcgccaggctggagtgagtgaggatctcggtcactgca
1231 agctccgctcccagggtccacgccattctcctgcctcagcctcccaagtagctgggactacaggcacatgccaccacacccg
1313 actaacttttttgtattttcagtagagacgggggtttcaccatgttggccaggctgggtctggaactcctgacctcaggtgatc

3/11

FIG. 2B

1395 tgcctggtttcgatctcccaaagtgtgggattacaggcgtgagccaccgcacctggctgggagtttagaggttttctaattgcat
1477 tgcaggcagatagtgaataccagacacggggcagctgtgatctttattctccatcacccccacacagccctgcctggggcac
1559 acaaggacactcaatacatgcttttccgctggggcgggtgggtcaccctgtaatcccagcactttgggaggccaaggtggga
1641 ggatcacttgagccaggagttcaacaccagcctgggcaacatagttagaccctgtctctactaaaaatacaaaaattagcc
1723 aggcattggtgccacacacctgtgctctcagctactcaggaggtgaggcaggaggatcgcttgagcccagaaggtcaaggtt
1805 gcagtgaacctatgttcaggccgctgcactccagcctgggtgacagagcaagacctgtttataaatacataatgctttccaa
1887 gtgattaaaccgactccccctcaccctgcccaccatgggtccaaagaagcatttgtggagcaccttctgtgtgcccctagg
1969 tagctagatgcctggacgggggtcagaaggacctgacccgaccttgaactgttccacacagGATGCCAGGCCAAGGTGGAG
G C Q A K V E
2051 CAAGCGGTGGAGACAGAGCCGGAGCCCGAGCTGCCGCCAGCAGACCGAGTGGCAGAGCGGCCAGCGCTGGGAAGTGGCACTGG
Q A V E T E P E P E L R Q Q T E W Q S G Q R W E L A L
2133 GTCGCTTTTGGGATTACCTGCCGCTGGGTGCAGACACTGTCTGAGCAGGTGCAGGAGGAGCTGCTCAGCTCCAGGTCAACCA
G R F W D Y L R W V Q T L S E Q V Q E E L L S S Q V T Q
2215 GGAAGTGAAGgtgagtggtcccatcctggcccttgacctcctgggtggggtatatacctccccagggtccaggtttcattctg
E L R
2297 cccctgtcgctaagtcttggggggcctgggtctctgtggttcttagcttctcttcccatttctgactcctggctttagctc
2379 tctggaattctctctctcagctttgtctctctctcttcccttctgactcagctctctcacactcgtcctggctctgtctctgt
2461 ccttccctagctcttttatatagagacagagagatggggctcactgtgttgcccaggctgggtcttgaacttctgggtcaa
2543 gcgatcctccccgctcgccctcccaaagtgtgggattagaggcatgagcaccttgcccgccctcctagctccttcttcgtc
2625 tctgcctctgcctctgcctctgtctctgtcatctgtctctgtctctctctccttctctcgccctctgcccgttctctctcctc
2707 ttgggtctctctgggtcatccccatctcgcccgcccatcccagcccttctccccgcctccccactgtgagacacctccc

4/11

FIG. 2C

2789 gccctctcggccgcagGGCGCTGATGGACGAGACCATGAAGGAGTTGAAGGCCTACAAATCGGAACCTGGAGGAACAACCTGAC
A L M D E T M K E L K A Y K S E L E Q L T

2871 CCCGGTGGCGGAGGAGACGCGGGCACGGCTGTCCAAGGAGCTGCAGGCGGCGCAGGCCCGGCTGGGCGCGGACATGGAGGAC
P V A E E T R A R L S K E L Q A A Q A R L G A D M E D

2953 GTGTGCGGCGCCTGGTGCAGTACCGCGGCGAGGTGCAGGCCATGCTCGGCCAGAGCACCGAGGAGCTGCGGGTGGCGCCTCG
V C G R L V Q Y R G E V Q A M L G Q S T E E L R V R L

3035 CCTCCACCTGCGCAAGCTGCGTAAGCGGCTCCTCCGCGATGCCGATGACCTGCAGAAGCGCCTGGCAGTGTACCAGGCCGG
A S H L R K L R K R L L R D A D D L Q K R L A V Y Q A G

3117 GGGCGCGAGGGCGCGAGCGCGGCTCAGCGCCATCCGCGAGCGCCTGGGGCCCTGGTGAACAGGGCCGCGTGGCGGCC
A R E G A E R G L S A I R E R L G P L V E Q G R V R A

3199 GCCACTGTGGGCTCCCTGGCGCGCCAGCCGCTACAGGAGCGGGCCAGGCCCTGGGGCGAGCGGCTGCGCGCGGATGGAGG
A T V G S L A G Q P L Q E R A Q A W G E R L R A R M E

3281 AGATGGGCGAGCCGACCCGCGACCGCCTGGACGAGGTGAAGGAGCAGGTGGCGGAGGTGCGCGCCAAAGCTGGAGGAGCAGGC
E M G S R T R D R L D E V K E Q V A E V R A K L E E Q A

3363 CCAGCAGATACGCCTGCAGGCGGAGGCCTTCCAGGCCCGCCTCAAGAGCTGGTTGAGCCCTGGTGAAGACATGCAGCGC
Q Q I R L Q A E A F Q A R L K S W F E P L V E D M Q R

3445 CAGTGGGCGGGCTGGTGGAGAAGGTGCAGGCTGCCGTGGGCACCGCGCGCCCTGTGCCAGCGACAATCACTGAACGC
Q W A G L V E K V Q A A V G S A A P V P S D N H

3527 CGAAGCCTGCAGCCATGCGACCCACGCCACCCCGTGCCTCCTGCCTCCGCGCAGCCTGCAGCGGGAGACCCCTGTCCCCGCC

3609 CCAGCCGTCTCTCTGGGGTGGACCTAGTTTAATAAAGATTACCAAGTTTCACGCActctgctggcctccccctgtgatttc

3691 ctctaagccccagcctcagttttctctttctgcccacatactgccacacaattctcagccccctctctccatctgtgtctgt

3373 gtgtatctttctctctgccccttttttttttttt

5/11

FIG. 3A

1 ctaaagaaga gactggtgg gagacaggg cgggggaagg gggaggggag tgaagtagtc
61 tccctggaat gctggtggtg ggggagggcag tctccttgggt ggaggagtcc cagcgtccct
121 cccctccct cctctgcca cacaatggac aatggcaact gccacacac tccatggag
181 gggaagggga tgagtgcagg gaaccccgac ccacccggg agacctgcaa gcctgcagca
241 ctccctccc gccccactg aacccttgac ccctgccctg cacgccccgc agcttgctgt
301 ttgcccactc ctatttgccc agtcccaggg acagagctga tccttgaaact cttaagtctc
361 acattgccag gaccagtga cagcaacagg gccagggctg ggcttatcag cctcccagcc
421 cagaccctgg ctgcagacat aaataggccc tgcaagagct ggctgcttag agactgcgag
481 aaggagggtg gtctgtgctg ctgccccggc actctggctc ccagctcaa ggttcaggcc
541 ttgccccagg ccgggcctct gggtaacctg ggtcttctcc cgctctgtgc ccttctctc
601 acctggctgc aatgagtggg ggagcacggg gctctgcat gctgaaggca cccactcag
661 ccaggccctt cttctctcc aggtccccc cggcccttca ggatgaaagc tgcggtgctg
721 accttgccg tgctcttctt gacgggtagg tgtccccctaa ctaggagcc aaccatcggg
781 gggcttctc cctaaatccc cgtggcccac cctcctgggc agaggcagca ggttctctc
841 tggcccccct tccccacct ccaagcttgg ccttctggct cagatctcag ccacagctg
901 gcctgatctg ggtctcccct cccacctca gggagccagg ctcggcattt ctggcagcaa
961 gatgaacccc ccagagccc ctgggatcga gtgaaggacc tggccactgt gtacgtggat
1021 gtgctcaaag acagcggcag agactatgtg tcccagttg aaggctccgc cttgggaaaa
1081 cagctaaagt aaggaccag cctgggggtg agggcagggg cagggggcag aggcctgtgg
1141 gatgatgtg aagccagact ggccgagtcc tcacctataa tctgatgagc tgggccccac
1201 agatggtctg gatggagaaa ccggaatgga tctccaggca gggtcacagc ccatgtcccc
1261 tgcaaaaggac agaccagggc tgcccgatgc gtgatcacag agccacattg tgcctgcaag
1321 tgtagcaagc cccttctccct tcttcaccac ctctctgct cctgcccagc aagactgtgg
1381 gctgtctctc gagaggagaa tgcgctggag gcatagaagc gaggtccttc aagggccac

6/11

FIG. 3B

1441 tttggagacc aacgtaactg ggcaccagtc ccagctctgt ctcttttta gctcctctct
1501 gtgcctcggc ccagctgcac aacggggcat ggccctggcg ggaggggtg ttggttgaga
1561 gtgtactgga aatgctaggc cactgcacct ccgcgacag gtgtcaccca gggctcacc
1621 ctgataggct gggcgctgg gaggccagcc ctcaaccctt ctgtctcacc ctccagccta
1681 aagctccttg acaactggga cagctgacc tcaccttca gcaagctgcg cgaacagctc
1741 gcccctgtga ccaggagtt ctgggataac ctggaaaagg agacagaggg cctgaggcaa
1801 gagatgagca aggatctgga ggaggtgaag gccaaaggcg agccctacct ggacgacttc
1861 cagaagaagt ggcaggagga gatggagctc taccgccaga agtgggagcc gctgcgcgca
1921 gagctccaag agggcgcgcg ccagaagctg cacgagctgc aagagaagct gagccactg
1981 ggcgaggaga tgcgcgaccg cgcgcgcgcc catgtggacg cgctgcgcac gcatctggcc
2041 ccctacagcg acgagctgcg ccagcgcttg gccgcgcgcc ttgaggctct caaggagaac
2101 ggcggcgcca gactggcga gtaccagcc aaggccaccg agcatctgag cagctcagc
2161 gagaaggcca agcccgcgct cgaggacctc cgccaaaggc tgctgcccgt gctggagagc
2221 ttcaaggcca gcttcctgag cgctctcgag gactacacta agaagctcaa caccagtga
2281 ggcgcccgcg gccgcccccc ttcccggcgc tcagaaataa cgtttccaaa tggggaagca
2341 gcttcttctt ttgggagaaa tagagggggg tgcggggaca tccgggggag cccgggaggg
2401 gcctttggcc ctggagcagg gacttcctgc cggatctcaa caactccgtg ccagactgg
2461 acgtcttagg gccaaagatcg acgttgagg acctgctgga cgcntggctg cttacgagtg
2521 agggagtaga gtctgcctta gcaaggctca agtagaaaagg aagtcacagc ggacnaggca
2581 aagcdacaga caatccaagg ccagggtgccc tgaagggggc tcaaaacaagg cctgcagcct
2641 gtctgaggcg ggcaggaaa cagcggttgc tttagctggg agcagtcggg tccccgtcc
2701 ccagaggtgt gtccgtatag agccttctcc agcccagccg ctgtcagcgg gcgggacgga
2761 gcggggcgcc tcaggagacc agccactggg attgggggtt ggtccccggg gcaagtgaag
2821 cgcttgaggt ttgcgcctgt cctcctttac taattcaaaa acctctcaa cagacacttc

7/11

FIG. 3C

2881 ccttttcttc tcacaaggcc agtatccccc tcccactact cccatcccg ccagaaacag
 2941 ccgcggttc ctcaggcaca gcagtgaag ccagtcctcc acccctgcg gctccatgcc
 3001 atgccacccc ctctttctgc cagccctggc agaagctggc ctgagtaaga aaattcacca
 3061 ccacctcttg caggtacatt tttatttcca agatgctctc atatctgtgc tctcactgca
 3121 tcctcccttc ccacatcct ggctagattg ccatcagacg cagagcatgg atgaggacac
 3181 tgaagcctgg acctgtgacg tcgcttgccc agtgaacagc aggatggct aggcagcgct
 3241 ttttagacc tgcacccctg gccatccatg attattgaaa agagtgtgcg ggtcgggtgc
 3301 ggtggctcaa gcctgtaatc ccagcacttt gggaggctga ggtgggcgta tcacttcagg
 3361 gccaggagtt tgagaccagc ctggccaata tggtgaaacc ctgtctctac taaaaatata
 3421 aaaaaaaat cagctggcat ggtggcttgc accgtaatc ccagctacta ggaaggctga
 3481 ggcaggagaa tcgcttgaac ctgggaggca gaggtcacag tgagccgaaa tcatgccact
 3541 gcactccagc ctgggcgacg gagcaagact ccagctaaaa aaaaaaaaag
 3601 agtgtgtggc ctggcactca agttcacatg ggtgtgcagg catgcctgtg tattctcaca
 3661 tgacctccct gctcacggtc cctccttgca ctcatgtctg aatgtcccg ctgcacgcac
 3721 atggcttcac agatctggca gtgccttccc taccctctct ctgcagggcc ttttgcccc
 3781 tcatgcaggc ccctggataa tcggcccccatt cccatgtcc ccatctccag tgtatcttag
 3841 ctaccctagg taaaggagtg ggcttttttag ttctaacct tccagagcta caacagcagt
 3901 catccagcca ggtctgggtg ggaacathtt ctagatacgg gtgctgagat ctctcagccc
 3961 agagagaagc cctggggaat tttcagagag aaagcagtct ccaggtgggg ctggatgtac
 4021 tgatgccact gagatctgta aaggagtccc taacacctga cataggagtg acaaaactgt
 4081 tttctgcacc aactgagcag aatacacgca gctgacctgg gctcaaggtc tggccctgcc
 4141 acgtgctggc tctgtgatgc tggccaagtg ccttcgcctc tccggggccac agttttttga
 4201 tctgaagagt ggagccctac tcaagccatc tgcagctctc gggctctctg acctgacatc
 4261 tttcgggtgg tggggacaca aaggaagcag cctctattgg gagacctgt gcttctttt

8/11

FIG. 3D

4321 ggtcccagga cactgcccc caccactcca gtccggtccc aaggcccag tcagctcaac
4381 tgtaatcatg acaacattga tcaagcatct ttacgtgcag gtgctgtgcc aaacggttcg
4441 aacgctctct cattcaatc tcacggcaaa cctacggtgg aggggtacg gttgtatcca
4501 ctttacatgt aagaaactga ggctgatata aagtgttga gccagaata gtgcctcgtt
4561 gcatcttact ccaacctcta gccatccgg cctcctccct tcacgtgcgc ctaagagggc
4621 taggtggcct ggatagggga ggtcagctcc acagttttga gtaaacacac acagtctcaa
4681 ctctgatgac acttaagtgc caggcatagt ggctggcatg gggcacacac tcaagtcatg
4741 ttgtgcagca cctaacagtt tatcaaatga tcagcaaaact tattgtcctg tttgaccttc
4801 cgacaaaagc tgtcaaggaa ggcaggggtac ggaggggtgat tctacctta gagatgaaga
4861 aactgaggcc cagagactag ccagctacc agaaggtgga tagagcgtg gcctccatgc
4921 ctgcctgacc tggagtctgt ccagtgtccc ccacagaaac agcctcggcc ctttcccatg
4981 ccagagacaca gatggcacac ttgcgacggc cactcatag cagcttcttg tccagcttta
5041 ttgggaggcc agcatgcctg gagggggggc aggcattgag tggggtagga ggcactgag
5101 aatactgtcc ctttaagca acctacaggg gaagccctgg agattgcagg acccaaggag
5161 ctggcaggat ggataggcag gtggacttgg ggtattgagg tctcaggcag ccacggctga
5221 agttggtctg acctcagggt ccaaatccca gaactcagag aacttgtcct taacggtgct
5281 ccagtagtct ttcagggaac tgaagccatc ggtcacccag cccctaaatc agtcagggga
5341 agcaaacagag cagggcatga gaactcctct gtaggcaacc atgggaccca caccatgtc
5401 cccactggac gacaccagtc aggaccacac caccctctca acttcactgg acgacagccc
5461 tgagacctca ggcagggaatc cccacagct gccacctgg gagaagagat atccttgcag
5521 gaacccccag acaagtcaaa ccctgccatc tccaggtcac ctcatgtt tatgccccctg
5581 ggcctgaggc acaaagtga aggtgggga gatgtgaaag gtcaaggctg tcattgtttt
5641 ctgtgcaaac agcaccacct ggagttgcac aacctggtgg ctctgagcag ggtaggacag
5701 agggaggcag cctctcattt ggaggattg attagtgtg tgatctggg

9/11

FIG. 3E

5761 caggtcacct aatcgctctg agcctcaatt ttctcatctg caaagtgaga aaataacacc
 5821 taccctaaag ccggttcttg ggactaagaa tgtttatgaa cacctctgct atgccagcta
 5881 atgccagtca ggtgaggtg gagaaggag taggggagag gaggtactg atgggcaggg
 5941 gcaggatggg agaggaaggc cacacttctg ctcacactgg cccaggtct tagaacaagg
 6001 cagaagcact cacgggcttg aattgggtca ggtggggcct cccaaggcaa accctagcc
 6061 cagggtcccc aacccccaaag tcatagatag gtactggtcc atggactgtt aggagccggg
 6121 ctgcacagca cgaagtgagt ggtgggtgag tgagcaatac gcctgagctc cgcctcctgt
 6181 cagatcaagt ggccttacct tctcatagga gtgtgaacct tattgtgaac tgcacatgcc
 6241 agggatctag gtggagggt ccttatgaga atctaatac tgatgatctg aggtggaaca
 6301 gcttcatccc aaaccatcc ccgctggccc atggaaaaat tgtccaccac aaaactggtc
 6361 cctggtgcca aaaggctga ggaccactgc tctagccaga ccttcagaaa aggaaaatgg
 6421 ggccaggcgc agtggctcgt gcctgtaatc ccagcacttt gggaggccga ggcaggagga
 6481 tcccctgagg tcaggaattc aagaccaacc tggccaacat ggtgaaaccc catctctgct
 6541 aaaaatacaa aaattagctg ggtgtggttg cgcgtgcctg taatcccagc tacttgggag
 6601 gctgaggcag gagaatgggt tgaaccggg agacggaggt tgcatgtgag cgagatggca
 6661 cactgcact ccagcctggg tgacagaggg agactccatt aaaaaaaa gaaagaaaa
 6721 gaaagagaga aaggaaagaa gggaaagaaa gaaaaaggaa agaaagaaaa
 6781 aaagaaaaa aagaaagaaa ggaagaaaaa gaaagggaaa gaaagggaaa
 6841 aagaaagaaa gaaagaaa aagaaaaata gaaagagaaa gaaagggaaa
 6901 atggaaatga gggctaaac ggcgcggccc taggactgct ccgggagaaa ggaggacct
 6961 gggaggaatg cagtctcaac tctgtcatct ctcccgcagc agcctgacaa aggccctgtg
 7021 aggaaagagg aggtgaaga gcacaggccc aggggagggt ggagcacctc cattccattg
 7081 ttgggatctc accagggcag gggcgggttg gaatggaggc agctggcagg ggggatggg
 7141 agggaggcca gcgggtgtac ctggcctgct gggccacctg ggactcctgc acgctgctca

10/11

FIG. 3F

7201 gtgcatacctt ggcggtcttg gtggcgtgct tcatgtagcc ctgcatgaag ctgagaaagg
7261 aggcatacctc ggcctcttgaa gctcctgagg aaagagcagg gctgagtggg gtggatcggc
7321 ctctggacga gccctggggc tctgcttga ccaccattg gactgggac cccaagtgc
7381 ctccaccctg ccccagccc agtcccacca agtgcttacg ggcagaggcc aggagcgcca
7441 ggagggaac aacaaggagt acccggggt gcattggacc tctgttcctg caaggaagt
7501 tcttgtgagg ggcacccag gtcccatgc ctctggacc ctccctggg aggtggcgtg
7561 gccctaaagg tagaacctta gctgggtctg ccagaaggag taggggcccg ctctgctca
7621 tacgggtctc cagaaggggg actggtgagc ggcgagggat cgaggccaa aggaggtggg
7681 tgggatggag cagaaaacc accagactga acatcaagg actgcgtctt ggaactgatc
7741 ccgtccagtc cagccaacat gctgtgtgtc ttgtgggtgat ttctggccct ctccaggcc
7801 tcagtttccc tgtctgggt aggaactggc tgtctaaagg ttctccagc cctaagcctg
7861 aagaatgagg ggggaacctg cacttgagc cacttccagc ccacccctt atgtagcttt
7921 gggcaagtga cactcccgg gcctccatgt tcttcaggtt atgatgagg gtggggggca
7981 cccgtccagc tccgaggctt ccttagctct agcaagtgt tctccaggct tgctggctgg
8041 gctgggcagg gagctcctc tgcctctct catcctcctc ccctcctt tccctcccc
8101 agagggcat accctggagc gctgcctcta gggatgaact gagcagacag gcaggagggt
8161 tctgacctgt ttatatcat ctccaggga caggcactg aggaccagg gcgctgggca
8221 aaggtcacct gctgaccagt ggagatgagg gcctgaggca ggtgtccag atgcagcaag
8281 cgggcgggag agttggaaa tccctaggag actgagtcca cgctgtgtc ccgccagccc
8341 tgcagcccag atgagctcag gaactggggg tgggcctggg gacctcatt ccttcctagc
8401 tgactggctc ccagggaga ggctgggtgag aggggaaatg gcttggacat agggcagcgc
8461 tcgggcccac ctacgcctt cacactggaa ttbcaggccc ctccctccac cagccccaa
8521 ccctgaacac agcctggagt agaggggtga ggggcttctt cagacttgag acaagtggg
8581 tggcttgggc tggggggtgt ttggagtaaa ggcacagaag accgggcac agtggctcgc

11/11

FIG. 3G

8641 aaggccctggt tacaggggctg agctcacagc ccctcccagc acctccatct ctgggttttca
8701 atccaggcag gcgagtgctg ctaccccagc agccccccac ccgccccac cctgtgtgcc
8761 ccaccgccgc ctccccctga gtgtagggca ggggttggtg gagaagcgca aggccgctca
8821 gagccccgagg cctttgcccc tccctccacc aggtcccta tttgcccctc tggaccact
8881 gataacatcc ctggggggagg agccccacc aggactgatt ctctcgttca ccttcctggg
8941 ttcctcagtg gggctggggg aagagc

SEQUENCE LISTING

<110> 1. Steer, Clifford J.
 2. Kren, Betsy T.
 3. Bandyopadhyay, Paramita
 4. Roy-Chowdhury, Jayanta

<120> Methods and Compounds for the Genetic
 Treatment of Hyperlipidemia

<130> 7991-033-228

<150> 60/074,497

<151> 1998-02-12

<150> 09/108,006

<151> 1998-06-30

<160> 59

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 4563

<212> PRT

<213> Homo Sapiens

<400> 1

Met	Asp	Pro	Pro	Arg	Pro	Ala	Leu	Leu	Ala	Leu	Leu	Ala	Leu	Pro	Ala
1				5					10					15	
Leu	Leu	Leu	Leu	Leu	Leu	Ala	Gly	Ala	Arg	Ala	Glu	Glu	Glu	Met	Leu
			20				25						30		
Glu	Asn	Val	Ser	Leu	Val	Cys	Pro	Lys	Asp	Ala	Thr	Arg	Phe	Lys	His
		35				40						45			
Leu	Arg	Lys	Tyr	Thr	Tyr	Asn	Tyr	Glu	Ala	Glu	Ser	Ser	Ser	Gly	Val
	50				55						60				
Pro	Gly	Thr	Ala	Asp	Ser	Arg	Ser	Ala	Thr	Arg	Ile	Asn	Cys	Lys	Val
65				70					75					80	
Glu	Leu	Glu	Val	Pro	Gln	Leu	Cys	Ser	Phe	Ile	Leu	Lys	Thr	Ser	Gln
			85						90					95	
Cys	Ile	Leu	Lys	Glu	Val	Tyr	Gly	Phe	Asn	Pro	Glu	Gly	Lys	Ala	Leu
		100					105						110		
Leu	Lys	Lys	Thr	Lys	Asn	Ser	Glu	Glu	Phe	Ala	Ala	Ala	Met	Ser	Arg
		115					120					125			
Tyr	Glu	Leu	Lys	Leu	Ala	Ile	Pro	Glu	Gly	Lys	Gln	Val	Phe	Leu	Tyr
	130				135						140				
Pro	Glu	Lys	Asp	Glu	Pro	Thr	Tyr	Ile	Leu	Asn	Ile	Lys	Arg	Gly	Ile
145				150					155					160	
Ile	Ser	Ala	Leu	Leu	Val	Pro	Pro	Glu	Thr	Glu	Glu	Ala	Lys	Gln	Val
			165						170					175	
Leu	Phe	Leu	Asp	Thr	Val	Tyr	Gly	Asn	Cys	Ser	Thr	His	Phe	Thr	Val
		180						185					190		
Lys	Thr	Arg	Lys	Gly	Asn	Val	Ala	Thr	Glu	Ile	Ser	Thr	Glu	Arg	Asp

2/11

FIG. 2A

1 cctatccctgggggagggggcgggacagggggagccctataattggacaagtctgggatccttgagtcctACTCAGCCCCAG
83 CGGAGGTGAAGGACGTCTTCCCCAGGAGCGGgtgagaagcgcagtcgggggacggggatgagctcaggggcctctagaaa
165 gagctgggaccctgggaagccctggcctccaggtagtctcaggagagctactcggggtcgggcttggggagaggaggagcgg
247 ggggtaggcaagcagcaggggactggacctgggaagggtgggcagcagagacgacccgacccgctagaagggtggggtgggg
329 agagcagctggactgggatgtaagccatagcaggactccacgagttgtcactatcattatcgagcacctactgggtgtcccc
411 agtgtcctcagatctccataactggggagccaggggcagcgacacggtagctagccgtcgattggagaactttaaaatgagg
493 actgaattagctcataaatggaaacagcgcttaactgtgaggttggagcttagaatgtgaagggagaatgaggaaatgcgag
575 actgggactgagatggaaacggcggtggggagggggtgggggatggaatttgaacccgggagaggaagatggaattttct
657 atggaggccgacctggggatggggagataagagaagaccaggagggttaaatagggaatgggttgggggagggttggtaa
739 atgtgctgggattaggctgttcagataatgcaacaaggcttggaggctaacctggggtgaggccgggttgggggagctgg
821 ggggtgggaggagtccctcactggcggttgattgacagttttctcttccccagACTGGCCAATCAGGCAGGAAGATGAAGGT
M K V
903 TCTGTGGGCTGCGTTGCTGTGCACATTCTGGCAGgtatgggggcggggcttgctcggttccccccgctcctccccctctca
L W A A L L V T F L A
985 tcctcacctcaacctcctggccccattcagacagacctgggccccctcttctgaggcttctgtgctgcttcctggctctga
1067 acagcgatttgacgctctctgggcctcggtttccccatccttgagataggagttagaagttgttttgttgttgttgtt
1149 tgttgttgttttgtttttttagatgaagtctcgctctgtcgccaggtggagtgcagtgggggatctcggtcactgca
1231 agctccgctcccagggtccacgccatttctctgctcagcctcccaagtagctgggactacaggcacatgccaccacacccg
1313 actaacttttttgtattttcagtagagacggggttcaccatgttggccagggtggtctggaactcctgacctcaggtgatc

Lys Ser Val Ser Ile Pro Ser Leu Asp Pro Ala Ser Ala Lys Ile Glu
 645 650 655
 Gly Asn Leu Ile Phe Asp Pro Asn Asn Tyr Leu Pro Lys Glu Ser Met
 660 665 670
 Leu Lys Thr Thr Leu Thr Ala Phe Gly Phe Ala Ser Ala Asp Leu Ile
 675 680 685
 Glu Ile Gly Leu Glu Gly Lys Gly Phe Glu Pro Thr Leu Glu Ala Leu
 690 695 700
 Phe Gly Lys Gln Gly Phe Phe Pro Asp Ser Val Asn Lys Ala Leu Tyr
 705 710 715 720
 Trp Val Asn Gly Gln Val Pro Asp Gly Val Ser Lys Val Leu Val Asp
 725 730 735
 His Phe Gly Tyr Thr Lys Asp Asp Lys His Glu Gln Asp Met Val Asn
 740 745 750
 Gly Ile Met Leu Ser Val Glu Lys Leu Ile Lys Asp Leu Lys Ser Lys
 755 760 765
 Glu Val Pro Glu Ala Arg Ala Tyr Leu Arg Ile Leu Gly Glu Glu Leu
 770 775 780
 Gly Phe Ala Ser Leu His Asp Leu Gln Leu Leu Gly Lys Leu Leu Leu
 785 790 795 800
 Met Gly Ala Arg Thr Leu Gln Gly Ile Pro Gln Met Ile Gly Glu Val
 805 810 815
 Ile Arg Lys Gly Ser Lys Asn Asp Phe Phe Leu His Tyr Ile Phe Met
 820 825 830
 Glu Asn Ala Phe Glu Leu Pro Thr Gly Ala Gly Leu Gln Leu Gln Ile
 835 840 845
 Ser Ser Ser Gly Val Ile Ala Pro Gly Ala Lys Ala Gly Val Lys Leu
 850 855 860
 Glu Val Ala Asn Met Gln Ala Glu Leu Val Ala Lys Pro Ser Val Ser
 865 870 875 880
 Val Glu Phe Val Thr Asn Met Gly Ile Ile Ile Pro Asp Phe Ala Arg
 885 890 895
 Ser Gly Val Gln Met Asn Thr Asn Phe Phe His Glu Ser Gly Leu Glu
 900 905 910
 Ala His Val Ala Leu Lys Pro Gly Lys Leu Lys Phe Ile Ile Pro Ser
 915 920 925
 Pro Lys Arg Pro Val Lys Leu Leu Ser Gly Gly Asn Thr Leu His Leu
 930 935 940
 Val Ser Thr Thr Lys Thr Glu Val Ile Pro Pro Leu Ile Glu Asn Arg
 945 950 955 960
 Gln Ser Trp Ser Val Cys Lys Gln Val Phe Pro Gly Leu Asn Tyr Cys
 965 970 975
 Thr Ser Gly Ala Tyr Ser Asn Ala Ser Ser Thr Asp Ser Ala Ser Tyr
 980 985 990
 Tyr Pro Leu Thr Gly Asp Thr Arg Leu Glu Leu Glu Leu Arg Pro Thr
 995 1000 1005
 Gly Glu Ile Glu Gln Tyr Ser Val Ser Ala Thr Tyr Glu Leu Gln Arg
 1010 1015 1020
 Glu Asp Arg Ala Leu Val Asp Thr Leu Lys Phe Val Thr Gln Ala Glu
 1025 1030 1035 1040
 Gly Ala Lys Gln Thr Glu Ala Thr Met Thr Phe Lys Tyr Asn Arg Gln
 1045 1050 1055
 Ser Met Thr Leu Ser Ser Glu Val Gln Ile Pro Asp Phe Asp Val Asp
 1060 1065 1070
 Leu Gly Thr Ile Leu Arg Val Asn Asp Glu Ser Thr Glu Gly Lys Thr

1075	1080	1085
Ser Tyr Arg Leu Thr Leu Asp Ile Gln Asn Lys Lys Ile Thr Glu Val		
1090	1095	1100
Ala Leu Met Gly His Leu Ser Cys Asp Thr Lys Glu Glu Arg Lys Ile		
1105	1110	1115
Lys Gly Val Ile Ser Ile Pro Arg Leu Gln Ala Glu Ala Arg Ser Glu		
	1125	1130
Ile Leu Ala His Trp Ser Pro Ala Lys Leu Leu Leu Gln Met Asp Ser		
	1140	1145
Ser Ala Thr Ala Tyr Gly Ser Thr Val Ser Lys Arg Val Ala Trp His		
	1155	1160
Tyr Asp Glu Glu Lys Ile Glu Phe Glu Trp Asn Thr Gly Thr Asn Val		
	1170	1175
Asp Thr Lys Lys Met Thr Ser Asn Phe Pro Val Asp Leu Ser Asp Tyr		
1185	1190	1195
Pro Lys Ser Leu His Met Tyr Ala Asn Arg Leu Leu Asp His Arg Val		
	1205	1210
Pro Gln Thr Asp Met Thr Phe Arg His Val Gly Ser Lys Leu Ile Val		
	1220	1225
Ala Met Ser Ser Trp Leu Gln Lys Ala Ser Gly Ser Leu Pro Tyr Thr		
	1235	1240
Gln Thr Leu Gln Asp His Leu Asn Ser Leu Lys Glu Phe Asn Leu Gln		
	1250	1255
Asn Met Gly Leu Pro Asp Phe His Ile Pro Glu Asn Leu Phe Leu Lys		
1265	1270	1275
Ser Asp Gly Arg Val Lys Tyr Thr Leu Asn Lys Asn Ser Leu Lys Ile		
	1285	1290
Glu Ile Pro Leu Pro Phe Gly Gly Lys Ser Ser Arg Asp Leu Lys Met		
	1300	1305
Leu Glu Thr Val Arg Thr Pro Ala Leu His Phe Lys Ser Val Gly Phe		
	1315	1320
His Leu Pro Ser Arg Glu Phe Gln Val Pro Thr Phe Thr Ile Pro Lys		
	1330	1335
Leu Tyr Gln Leu Gln Val Pro Leu Leu Gly Val Leu Asp Leu Ser Thr		
1345	1350	1355
Asn Val Tyr Ser Asn Leu Tyr Asn Trp Ser Ala Ser Tyr Ser Gly Gly		
	1365	1370
Asn Thr Ser Thr Asp His Phe Ser Leu Arg Ala Arg Tyr His Met Lys		
	1380	1385
Ala Asp Ser Val Val Asp Leu Leu Ser Tyr Asn Val Gln Gly Ser Gly		
	1395	1400
Glu Thr Thr Tyr Asp His Lys Asn Thr Phe Thr Leu Ser Cys Asp Gly		
	1410	1415
Ser Leu Arg His Lys Phe Leu Asp Ser Asn Ile Lys Phe Ser His Val		
1425	1430	1435
Glu Lys Leu Gly Asn Asn Pro Val Ser Lys Gly Leu Leu Ile Phe Asp		
	1445	1450
Ala Ser Ser Ser Trp Gly Pro Gln Met Ser Ala Ser Val His Leu Asp		
	1460	1465
Ser Lys Lys Lys Gln His Leu Phe Val Lys Glu Val Lys Ile Asp Gly		
	1475	1480
Gln Phe Arg Val Ser Ser Phe Tyr Ala Lys Gly Thr Tyr Gly Leu Ser		
	1490	1495
Cys Gln Arg Asp Pro Asn Thr Gly Arg Leu Asn Gly Glu Ser Asn Leu		
1505	1510	1515
		1520

Arg Phe Asn Ser Ser Tyr Leu Gln Gly Thr Asn Gln Ile Thr Gly Arg
 1525 1530 1535
 Tyr Glu Asp Gly Thr Leu Ser Leu Thr Ser Thr Ser Asp Leu Gln Ser
 1540 1545 1550
 Gly Ile Ile Lys Asn Thr Ala Ser Leu Lys Tyr Glu Asn Tyr Glu Leu
 1555 1560 1565
 Thr Leu Lys Ser Asp Thr Asn Gly Lys Tyr Lys Asn Phe Ala Thr Ser
 1570 1575 1580
 Asn Lys Met Asp Met Thr Phe Ser Lys Gln Asn Ala Leu Leu Arg Ser
 1585 1590 1595 1600
 Glu Tyr Gln Ala Asp Tyr Glu Ser Leu Arg Phe Phe Ser Leu Leu Ser
 1605 1610 1615
 Gly Ser Leu Asn Ser His Gly Leu Glu Leu Asn Ala Asp Ile Leu Gly
 1620 1625 1630
 Thr Asp Lys Ile Asn Ser Gly Ala His Lys Ala Thr Leu Arg Ile Gly
 1635 1640 1645
 Gln Asp Gly Ile Ser Thr Ser Ala Thr Thr Asn Leu Lys Cys Ser Leu
 1650 1655 1660
 Leu Val Leu Glu Asn Glu Leu Asn Ala Glu Leu Gly Leu Ser Gly Ala
 1665 1670 1675 1680
 Ser Met Lys Leu Thr Thr Asn Gly Arg Phe Arg Glu His Asn Ala Lys
 1685 1690 1695
 Phe Ser Leu Asp Gly Lys Ala Ala Leu Thr Glu Leu Ser Leu Gly Ser
 1700 1705 1710
 Ala Tyr Gln Ala Met Ile Leu Gly Val Asp Ser Lys Asn Ile Phe Asn
 1715 1720 1725
 Phe Lys Val Ser Gln Glu Gly Leu Lys Leu Ser Asn Asp Met Met Gly
 1730 1735 1740
 Ser Tyr Ala Glu Met Lys Phe Asp His Thr Asn Ser Leu Asn Ile Ala
 1745 1750 1755 1760
 Gly Leu Ser Leu Asp Phe Ser Ser Lys Leu Asp Asn Ile Tyr Ser Ser
 1765 1770 1775
 Asp Lys Phe Tyr Lys Gln Thr Val Asn Leu Gln Leu Gln Pro Tyr Ser
 1780 1785 1790
 Leu Val Thr Thr Leu Asn Ser Asp Leu Lys Tyr Asn Ala Leu Asp Leu
 1795 1800 1805
 Thr Asn Asn Gly Lys Leu Arg Leu Glu Pro Leu Lys Leu His Val Ala
 1810 1815 1820
 Gly Asn Leu Lys Gly Ala Tyr Gln Asn Asn Glu Ile Lys His Ile Tyr
 1825 1830 1835 1840
 Ala Ile Ser Ser Ala Ala Leu Ser Ala Ser Tyr Lys Ala Asp Thr Val
 1845 1850 1855
 Ala Lys Val Gln Gly Val Glu Phe Ser His Arg Leu Asn Thr Asp Ile
 1860 1865 1870
 Ala Gly Leu Ala Ser Ala Ile Asp Met Ser Thr Asn Tyr Asn Ser Asp
 1875 1880 1885
 Ser Leu His Phe Ser Asn Val Phe Arg Ser Val Met Ala Pro Phe Thr
 1890 1895 1900
 Met Thr Ile Asp Ala His Thr Asn Gly Asn Gly Lys Leu Ala Leu Trp
 1905 1910 1915 1920
 Gly Glu His Thr Gly Gln Leu Tyr Ser Lys Phe Leu Leu Lys Ala Glu
 1925 1930 1935
 Pro Leu Ala Phe Thr Phe Ser His Asp Tyr Lys Gly Ser Thr Ser His
 1940 1945 1950
 His Leu Val Ser Arg Lys Ser Ile Ser Ala Ala Leu Glu His Lys Val

1955	1960	1965
Ser Ala Leu Leu Thr Pro Ala Glu Gln Thr Gly Thr Trp Lys Leu Lys		
1970	1975	1980
Thr Gln Phe Asn Asn Asn Glu Tyr Ser Gln Asp Leu Asp Ala Tyr Asn		
1985	1990	1995
Thr Lys Asp Lys Ile Gly Val Glu Leu Thr Gly Arg Thr Leu Ala Asp		2000
2005	2010	2015
Leu Thr Leu Leu Asp Ser Pro Ile Lys Val Pro Leu Leu Leu Ser Glu		
2020	2025	2030
Pro Ile Asn Ile Ile Asp Ala Leu Glu Met Arg Asp Ala Val Glu Lys		
2035	2040	2045
Pro Gln Glu Phe Thr Ile Val Ala Phe Val Lys Tyr Asp Lys Asn Gln		
2050	2055	2060
Asp Val His Ser Ile Asn Leu Pro Phe Phe Glu Thr Leu Gln Glu Tyr		
2065	2070	2075
Phe Glu Arg Asn Arg Gln Thr Ile Ile Val Val Leu Glu Asn Val Gln		2080
2085	2090	2095
Arg Asn Leu Lys His Ile Asn Ile Asp Gln Phe Val Arg Lys Tyr Arg		
2100	2105	2110
Ala Ala Leu Gly Lys Leu Pro Gln Gln Ala Asn Asp Tyr Leu Asn Ser		
2115	2120	2125
Phe Asn Trp Glu Arg Gln Val Ser His Ala Lys Glu Lys Leu Thr Ala		
2130	2135	2140
Leu Thr Lys Lys Tyr Arg Ile Thr Glu Asn Asp Ile Gln Ile Ala Leu		
2145	2150	2155
Asp Asp Ala Lys Ile Asn Phe Asn Glu Lys Leu Ser Gln Leu Gln Thr		
2165	2170	2175
Tyr Met Ile Gln Phe Asp Gln Tyr Ile Lys Asp Ser Tyr Asp Leu His		
2180	2185	2190
Asp Leu Lys Ile Ala Ile Ala Asn Ile Ile Asp Glu Ile Ile Glu Lys		
2195	2200	2205
Leu Lys Ser Leu Asp Glu His Tyr His Ile Arg Val Asn Leu Val Lys		
2210	2215	2220
Thr Ile His Asp Leu His Leu Phe Ile Glu Asn Ile Asp Phe Asn Lys		
2225	2230	2235
Ser Gly Ser Ser Thr Ala Ser Trp Ile Gln Asn Val Asp Thr Lys Tyr		
2245	2250	2255
Gln Ile Arg Ile Gln Ile Gln Glu Lys Leu Gln Gln Leu Lys Arg His		
2260	2265	2270
Ile Gln Asn Ile Asp Ile Gln His Leu Ala Gly Lys Leu Lys Gln His		
2275	2280	2285
Ile Glu Ala Ile Asp Val Arg Val Leu Leu Asp Gln Leu Gly Thr Thr		
2290	2295	2300
Ile Ser Phe Glu Arg Ile Asn Asp Val Leu Glu His Val Lys His Phe		
2305	2310	2315
Val Ile Asn Leu Ile Gly Asp Phe Glu Val Ala Glu Lys Ile Asn Ala		
2325	2330	2335
Phe Arg Ala Lys Val His Glu Leu Ile Glu Arg Tyr Glu Val Asp Gln		
2340	2345	2350
Gln Ile Gln Val Leu Met Asp Lys Leu Val Glu Leu Ala His Gln Tyr		
2355	2360	2365
Lys Leu Lys Glu Thr Ile Gln Lys Leu Ser Asn Val Leu Gln Gln Val		
2370	2375	2380
Lys Ile Lys Asp Tyr Phe Glu Lys Leu Val Gly Phe Ile Asp Asp Ala		
2385	2390	2395
		2400

Val Lys Lys Leu Asn Glu Leu Ser Phe Lys Thr Phe Ile Glu Asp Val
 2405 2410 2415
 Asn Lys Phe Leu Asp Met Leu Ile Lys Lys Leu Lys Ser Phe Asp Tyr
 2420 2425 2430
 His Gln Phe Val Asp Glu Thr Asn Asp Lys Ile Arg Glu Val Thr Gln
 2435 2440 2445
 Arg Leu Asn Gly Glu Ile Gln Ala Leu Glu Leu Pro Gln Lys Ala Glu
 2450 2455 2460
 Ala Leu Lys Leu Phe Leu Glu Glu Thr Lys Ala Thr Val Ala Val Tyr
 2465 2470 2475 2480
 Leu Glu Ser Leu Gln Asp Thr Lys Ile Thr Leu Ile Ile Asn Trp Leu
 2485 2490 2495
 Gln Glu Ala Leu Ser Ser Ala Ser Leu Ala His Met Lys Ala Lys Phe
 2500 2505 2510
 Arg Glu Thr Leu Glu Asp Thr Arg Asp Arg Met Tyr Gln Met Asp Ile
 2515 2520 2525
 Gln Gln Glu Leu Gln Arg Tyr Leu Ser Leu Val Gly Gln Val Tyr Ser
 2530 2535 2540
 Thr Leu Val Thr Tyr Ile Ser Asp Trp Trp Thr Leu Ala Ala Lys Asn
 2545 2550 2555 2560
 Leu Thr Asp Phe Ala Glu Gln Tyr Ser Ile Gln Asp Trp Ala Lys Arg
 2565 2570 2575
 Met Lys Ala Leu Val Glu Gln Gly Phe Thr Val Pro Glu Ile Lys Thr
 2580 2585 2590
 Ile Leu Gly Thr Met Pro Ala Phe Glu Val Ser Leu Gln Ala Leu Gln
 2595 2600 2605
 Lys Ala Thr Phe Gln Thr Pro Asp Phe Ile Val Pro Leu Thr Asp Leu
 2610 2615 2620
 Arg Ile Pro Ser Val Gln Ile Asn Phe Lys Asp Leu Lys Asn Ile Lys
 2625 2630 2635 2640
 Ile Pro Ser Arg Phe Ser Thr Pro Glu Phe Thr Ile Leu Asn Thr Phe
 2645 2650 2655
 His Ile Pro Ser Phe Thr Ile Asp Phe Val Glu Met Lys Val Lys Ile
 2660 2665 2670
 Ile Arg Thr Ile Asp Gln Met Leu Asn Ser Glu Leu Gln Trp Pro Val
 2675 2680 2685
 Pro Asp Ile Tyr Leu Arg Asp Leu Lys Val Glu Asp Ile Pro Leu Ala
 2690 2695 2700
 Arg Ile Thr Leu Pro Asp Phe Arg Leu Pro Glu Ile Ala Ile Pro Glu
 2705 2710 2715 2720
 Phe Ile Ile Pro Thr Leu Asn Leu Asn Asp Phe Gln Val Pro Asp Leu
 2725 2730 2735
 His Ile Pro Glu Phe Gln Leu Pro His Ile Ser His Thr Ile Glu Val
 2740 2745 2750
 Pro Thr Phe Gly Lys Leu Tyr Ser Ile Leu Lys Ile Gln Ser Pro Leu
 2755 2760 2765
 Phe Thr Leu Asp Ala Asn Ala Asp Ile Gly Asn Gly Thr Thr Ser Ala
 2770 2775 2780
 Asn Glu Ala Gly Ile Ala Ala Ser Ile Thr Ala Lys Gly Glu Ser Lys
 2785 2790 2795 2800
 Leu Glu Val Leu Asn Phe Asp Phe Gln Ala Asn Ala Gln Leu Ser Asn
 2805 2810 2815
 Pro Lys Ile Asn Pro Leu Ala Leu Lys Glu Ser Val Lys Phe Ser Ser
 2820 2825 2830
 Lys Tyr Leu Arg Thr Glu His Gly Ser Glu Met Leu Phe Phe Gly Asn

2835	2840	2845
Ala Ile Glu Gly Lys Ser Asn Thr Val	Ala Ser Leu His Thr Glu Lys	
2850	2855	2860
Asn Thr Leu Glu Leu Ser Asn Gly Val Ile	Val Lys Ile Asn Asn Gln	
2865	2870	2875
Leu Thr Leu Asp Ser Asn Thr Lys Tyr Phe	His Lys Leu Asn Ile Pro	2880
	2885	2890
Lys Leu Asp Phe Ser Ser Gln Ala Asp Leu	Arg Asn Glu Ile Lys Thr	2895
	2900	2905
Leu Leu Lys Ala Gly His Ile Ala Trp Thr	Ser Ser Gly Lys Gly Ser	2910
	2915	2920
Trp Lys Trp Ala Cys Pro Arg Phe Ser Asp	Glu Gly Thr His Glu Ser	2925
	2930	2935
Gln Ile Ser Phe Thr Ile Glu Gly Pro Leu	Thr Ser Phe Gly Leu Ser	2940
2945	2950	2955
Asn Lys Ile Asn Ser Lys His Leu Arg Val	Asn Gln Asn Leu Val Tyr	2960
	2965	2970
Glu Ser Gly Ser Leu Asn Phe Ser Lys Leu	Glu Ile Gln Ser Gln Val	2975
	2980	2985
Asp Ser Gln His Val Gly His Ser Val Leu	Thr Ala Lys Gly Met Ala	2990
	2995	3000
Leu Phe Gly Glu Gly Lys Ala Glu Phe Thr	Gly Arg His Asp Ala His	3005
	3010	3015
Leu Asn Gly Lys Val Ile Gly Thr Leu Lys	Asn Ser Leu Phe Phe Ser	3020
3025	3030	3035
Ala Gln Pro Phe Glu Ile Thr Ala Ser Thr	Asn Asn Glu Gly Asn Leu	3040
	3045	3050
Lys Val Arg Phe Pro Leu Arg Leu Thr Gly	Lys Ile Asp Phe Leu Asn	3055
	3060	3065
Asn Tyr Ala Leu Phe Leu Ser Pro Ser Ala	Gln Gln Ala Ser Trp Gln	3070
	3075	3080
Val Ser Ala Arg Phe Asn Gln Tyr Lys Tyr	Asn Gln Asn Phe Ser Ala	3085
	3090	3095
Gly Asn Asn Glu Asn Ile Met Glu Ala His	Val Gly Ile Asn Gly Glu	3100
3105	3110	3115
Ala Asn Leu Asp Phe Leu Asn Ile Pro Leu	Thr Ile Pro Glu Met Arg	3120
	3125	3130
Leu Pro Tyr Thr Ile Ile Thr Thr Pro Pro	Leu Lys Asp Phe Ser Leu	3135
	3140	3145
Trp Glu Lys Thr Gly Leu Lys Glu Phe Leu	Lys Thr Thr Lys Gln Ser	3150
	3155	3160
Phe Asp Leu Ser Val Lys Ala Gln Tyr Lys	Lys Asn Lys His Arg His	3165
	3170	3175
Ser Ile Thr Asn Pro Leu Ala Val Leu Cys	Glu Phe Ile Ser Gln Ser	3180
3185	3190	3195
Ile Lys Ser Phe Asp Arg His Phe Glu Lys	Asn Arg Asn Asn Ala Leu	3200
	3205	3210
Asp Phe Val Thr Lys Ser Tyr Asn Glu Thr	Lys Ile Lys Phe Asp Lys	3215
	3220	3225
Tyr Lys Ala Glu Lys Ser His Asp Glu Leu	Pro Arg Thr Phe Gln Ile	3230
	3235	3240
Pro Gly Tyr Thr Val Pro Val Val Asn Val	Glu Val Ser Pro Phe Thr	3245
	3250	3255
Ile Glu Met Ser Ala Phe Gly Tyr Val Phe	Pro Lys Ala Val Ser Met	3260
3265	3270	3275
		3280

Pro Ser Phe Ser Ile Leu Gly Ser Asp Val Arg Val Pro Ser Tyr Thr
 3285 3290 3295
 Leu Ile Leu Pro Ser Leu Glu Leu Pro Val Leu His Val Pro Arg Asn
 3300 3305 3310
 Leu Lys Leu Ser Leu Pro Asp Phe Lys Glu Leu Cys Thr Ile Ser His
 3315 3320 3325
 Ile Phe Ile Pro Ala Met Gly Asn Ile Thr Tyr Asp Phe Ser Phe Lys
 3330 3335 3340
 Ser Ser Val Ile Thr Leu Asn Thr Asn Ala Glu Leu Phe Asn Gln Ser
 3345 3350 3355 3360
 Asp Ile Val Ala His Leu Leu Ser Ser Ser Ser Val Ile Asp Ala
 3365 3370 3375
 Leu Gln Tyr Lys Leu Glu Gly Thr Thr Arg Leu Thr Arg Lys Arg Gly
 3380 3385 3390
 Leu Lys Leu Ala Thr Ala Leu Ser Leu Ser Asn Lys Phe Val Glu Gly
 3395 3400 3405
 Ser His Asn Ser Thr Val Ser Leu Thr Thr Lys Asn Met Glu Val Ser
 3410 3415 3420
 Val Ala Thr Thr Thr Lys Ala Gln Ile Pro Ile Leu Arg Met Asn Phe
 3425 3430 3435 3440
 Lys Gln Glu Leu Asn Gly Asn Thr Lys Ser Lys Pro Thr Val Ser Ser
 3445 3450 3455
 Ser Met Glu Phe Lys Tyr Asp Phe Asn Ser Ser Met Leu Tyr Ser Thr
 3460 3465 3470
 Ala Lys Gly Ala Val Asp His Lys Leu Ser Leu Glu Ser Leu Thr Ser
 3475 3480 3485
 Tyr Phe Ser Ile Glu Ser Ser Thr Lys Gly Asp Val Lys Gly Ser Val
 3490 3495 3500
 Leu Ser Arg Glu Tyr Ser Gly Thr Ile Ala Ser Glu Ala Asn Thr Tyr
 3505 3510 3515 3520
 Leu Asn Ser Lys Ser Thr Arg Ser Ser Val Lys Leu Gln Gly Thr Ser
 3525 3530 3535
 Lys Ile Asp Asp Ile Trp Asn Leu Glu Val Lys Glu Asn Phe Ala Gly
 3540 3545 3550
 Glu Ala Thr Leu Gln Arg Ile Tyr Ser Leu Trp Glu His Ser Thr Lys
 3555 3560 3565
 Asn His Leu Gln Leu Glu Gly Leu Phe Phe Thr Asn Gly Glu His Thr
 3570 3575 3580
 Ser Lys Ala Thr Leu Glu Leu Ser Pro Trp Gln Met Ser Ala Leu Val
 3585 3590 3595 3600
 Gln Val His Ala Ser Gln Pro Ser Ser Phe His Asp Phe Pro Asp Leu
 3605 3610 3615
 Gly Gln Glu Val Ala Leu Asn Ala Asn Thr Lys Asn Gln Lys Ile Arg
 3620 3625 3630
 Trp Lys Asn Glu Val Arg Ile His Ser Gly Ser Phe Gln Ser Gln Val
 3635 3640 3645
 Glu Leu Ser Asn Asp Gln Glu Lys Ala His Leu Asp Ile Ala Gly Ser
 3650 3655 3660
 Leu Glu Gly His Leu Arg Phe Leu Lys Asn Ile Ile Leu Pro Val Tyr
 3665 3670 3675 3680
 Asp Lys Ser Leu Trp Asp Phe Leu Lys Leu Asp Val Thr Thr Ser Ile
 3685 3690 3695
 Gly Arg Arg Gln His Leu Arg Val Ser Thr Ala Phe Val Tyr Thr Lys
 3700 3705 3710
 Asn Pro Asn Gly Tyr Ser Phe Ser Ile Pro Val Lys Val Leu Ala Asp

3715 3720 3725
 Lys Phe Ile Ile Pro Gly Leu Lys Leu Asn Asp Leu Asn Ser Val Leu
 3730 3735 3740
 Val Met Pro Thr Phe His Val Pro Phe Thr Asp Leu Gln Val Pro Ser
 3745 3750 3755 3760
 Cys Lys Leu Asp Phe Arg Glu Ile Gln Ile Tyr Lys Lys Leu Arg Thr
 3765 3770 3775
 Ser Ser Phe Ala Leu Asn Leu Pro Thr Leu Pro Glu Val Lys Phe Pro
 3780 3785 3790
 Glu Val Asp Val Leu Thr Lys Tyr Ser Gln Pro Glu Asp Ser Leu Ile
 3795 3800 3805
 Pro Phe Phe Glu Ile Thr Val Pro Glu Ser Gln Leu Thr Val Ser Gln
 3810 3815 3820
 Phe Thr Leu Pro Lys Ser Val Ser Asp Gly Ile Ala Ala Leu Asp Leu
 3825 3830 3835 3840
 Asn Ala Val Ala Asn Lys Ile Ala Asp Phe Glu Leu Pro Thr Ile Ile
 3845 3850 3855
 Val Pro Glu Gln Thr Ile Glu Ile Pro Ser Ile Lys Phe Ser Val Pro
 3860 3865 3870
 Ala Gly Ile Ala Ile Pro Ser Phe Gln Ala Leu Thr Ala Arg Phe Glu
 3875 3880 3885
 Val Asp Ser Pro Val Tyr Asn Ala Thr Trp Ser Ala Ser Leu Lys Asn
 3890 3895 3900
 Lys Ala Asp Tyr Val Glu Thr Val Leu Asp Ser Thr Cys Ser Ser Thr
 3905 3910 3915 3920
 Val Gln Phe Leu Glu Tyr Glu Leu Asn Val Leu Gly Thr His Lys Ile
 3925 3930 3935
 Glu Asp Gly Thr Leu Ala Ser Lys Thr Lys Gly Thr Phe Ala His Arg
 3940 3945 3950
 Asp Phe Ser Ala Glu Tyr Glu Glu Asp Gly Lys Tyr Glu Gly Leu Gln
 3955 3960 3965
 Glu Trp Glu Gly Lys Ala His Leu Asn Ile Lys Ser Pro Ala Phe Thr
 3970 3975 3980
 Asp Leu His Leu Arg Tyr Gln Lys Asp Lys Lys Gly Ile Ser Thr Ser
 3985 3990 3995 4000
 Ala Ala Ser Pro Ala Val Gly Thr Val Gly Met Asp Met Asp Glu Asp
 4005 4010 4015
 Asp Asp Phe Ser Lys Trp Asn Phe Tyr Tyr Ser Pro Gln Ser Ser Pro
 4020 4025 4030
 Asp Lys Lys Leu Thr Ile Phe Lys Thr Glu Leu Arg Val Arg Glu Ser
 4035 4040 4045
 Asp Glu Glu Thr Gln Ile Lys Val Asn Trp Glu Glu Glu Ala Ala Ser
 4050 4055 4060
 Gly Leu Leu Thr Ser Leu Lys Asp Asn Val Pro Lys Ala Thr Gly Val
 4065 4070 4075 4080
 Leu Tyr Asp Tyr Val Asn Lys Tyr His Trp Glu His Thr Gly Leu Thr
 4085 4090 4095
 Leu Arg Glu Val Ser Ser Lys Leu Arg Arg Asn Leu Gln Asp His Ala
 4100 4105 4110
 Glu Trp Val Tyr Gln Gly Ala Ile Arg Glu Ile Asp Asp Ile Asp Glu
 4115 4120 4125
 Arg Phe Gln Lys Gly Ala Ser Gly Thr Thr Gly Thr Tyr Gln Glu Trp
 4130 4135 4140
 Lys Asp Lys Ala Gln Asn Leu Tyr Gln Glu Leu Leu Thr Gln Glu Gly
 4145 4150 4155 4160

Gln Ala Ser Phe Gln Gly Leu Lys Asp Asn Val Phe Asp Gly Leu Val
 4165 4170 4175
 Arg Val Thr Gln Glu Phe His Met Lys Val Lys His Leu Ile Asp Ser
 4180 4185 4190
 Leu Ile Asp Phe Leu Asn Phe Pro Arg Phe Gln Phe Pro Gly Lys Pro
 4195 4200 4205
 Gly Ile Tyr Thr Arg Glu Glu Leu Cys Thr Met Phe Ile Arg Glu Val
 4210 4215 4220
 Gly Thr Val Leu Ser Gln Val Tyr Ser Lys Val His Asn Gly Ser Glu
 4225 4230 4235 4240
 Ile Leu Phe Ser Tyr Phe Gln Asp Leu Val Ile Thr Leu Pro Phe Glu
 4245 4250 4255
 Leu Arg Lys His Lys Leu Ile Asp Val Ile Ser Met Tyr Arg Glu Leu
 4260 4265 4270
 Leu Lys Asp Leu Ser Lys Glu Ala Gln Glu Val Phe Lys Ala Ile Gln
 4275 4280 4285
 Ser Leu Lys Thr Thr Glu Val Leu Arg Asn Leu Gln Asp Leu Leu Gln
 4290 4295 4300
 Phe Ile Phe Gln Leu Ile Glu Asp Asn Ile Lys Gln Leu Lys Glu Met
 4305 4310 4315 4320
 Lys Phe Thr Tyr Leu Ile Asn Tyr Ile Gln Asp Glu Ile Asn Thr Ile
 4325 4330 4335
 Phe Asn Asp Tyr Ile Pro Tyr Val Phe Lys Leu Leu Lys Glu Asn Leu
 4340 4345 4350
 Cys Leu Asn Leu His Lys Phe Asn Glu Phe Ile Gln Asn Glu Leu Gln
 4355 4360 4365
 Glu Ala Ser Gln Glu Leu Gln Gln Ile His Gln Tyr Ile Met Ala Leu
 4370 4375 4380
 Arg Glu Glu Tyr Phe Asp Pro Ser Ile Val Gly Trp Thr Val Lys Tyr
 4385 4390 4395 4400
 Tyr Glu Leu Glu Glu Lys Ile Val Ser Leu Ile Lys Asn Leu Leu Val
 4405 4410 4415
 Ala Leu Lys Asp Phe His Ser Glu Tyr Ile Val Ser Ala Ser Asn Phe
 4420 4425 4430
 Thr Ser Gln Leu Ser Ser Gln Val Glu Gln Phe Leu His Arg Asn Ile
 4435 4440 4445
 Gln Glu Tyr Leu Ser Ile Leu Thr Asp Pro Asp Gly Lys Gly Lys Glu
 4450 4455 4460
 Lys Ile Ala Glu Leu Ser Ala Thr Ala Gln Glu Ile Ile Lys Ser Gln
 4465 4470 4475 4480
 Ala Ile Ala Thr Lys Lys Ile Ile Ser Asp Tyr His Gln Gln Phe Arg
 4485 4490 4495
 Tyr Lys Leu Gln Asp Phe Ser Asp Gln Leu Ser Asp Tyr Tyr Glu Lys
 4500 4505 4510
 Phe Ile Ala Glu Ser Lys Arg Leu Ile Asp Leu Ser Ile Gln Asn Tyr
 4515 4520 4525
 His Thr Phe Leu Ile Tyr Ile Thr Glu Leu Leu Lys Lys Leu Gln Ser
 4530 4535 4540
 Thr Thr Val Met Asn Pro Tyr Met Lys Leu Ala Pro Gly Glu Leu Thr
 4545 4550 4555 4560
 Ile Ile Leu

<212> DNA

<213> Homo Sapiens

<400> 2

gaggagcccc	cccagccagc	cagggccgcg	aggccgaggc	cagggccgcg	cccaggagcc	60
gccccaccgc	agctggcgat	ggaccgcgcg	aggccgcgcg	tgctggcgct	gctggcgctg	120
cctgcgctgc	tgctgctgct	gctggcgggc	gccagggccg	aagaggaaat	gctggaaaat	180
gtcagcctgg	tctgtccaaa	agatgcgacc	cgattcaagc	acctccggaa	gtacacatac	240
aactatgagg	ctgagagttc	cagtggagtc	cctgggactg	ctgattcaag	aagtgccacc	300
aggatcaact	gcaagggttg	gctggagggt	ccccagctct	gcagcttcac	cctgaagacc	360
agccagtgc	tcctgaaaga	ggtgtatggc	ttcaaccctg	agggcaaagc	cttgctgaag	420
aaaaccaaga	actctgagga	gtttgctgca	gccatgtcca	ggtatgagct	caagctggcc	480
attccagaag	ggaagcaggt	tttccctttac	ccggagaaaag	atgaacctac	ttacatcctg	540
aacatcaaga	ggggcatcat	ttctgccctc	ctgggtccccc	cagagacaga	agaagccaag	600
caagtgttgt	ttctggatac	cgtgtatgga	aactgctcca	ctcactttac	cgtcaagacg	660
aggaagggca	atgtggcaac	agaaatatcc	actgaaagag	acctggggca	gtgtgatcgc	720
ttcaagccca	tcgcacacag	catcagccca	cttgctctca	tcaaaggcat	gacccgcccc	780
ttgtcaactc	tgatcagcag	cagccagtc	tgtcagtaca	cactggacgc	taagaggaag	840
catgtggcag	aagccatctg	caaggagcaa	cacctcttcc	tgcttttctc	ctacaagaat	900
aagtatggga	tggtagcaca	agtgcacacg	actttgaaac	ttgaagacac	accaaagatc	960
aacagccgct	tctttgggtga	aggtactaag	aagatggggc	tcgcatttga	gagcaccaaa	1020
tccacatcac	ctccaaagca	ggccgaagct	gttttgaaaga	ctgtccagga	actgaaaaaa	1080
ctaaccatct	ctgagcaaaa	tatccagaga	gctaactctc	tcaataagct	ggttactgag	1140
ctgagaggcc	tcagtgatga	agcagtcaca	tctctcttgc	cacagctgat	tgagggtgcc	1200
agccccatca	ctttacaagc	cttggttcag	tgtggacagc	ctcagtgtc	cactcacatc	1260
ctccagtggc	tgaaacgtgt	gcatgccaac	ccccttctga	tagatgtggt	cacctacctg	1320
gtggccctga	tcgccgagcc	ctcagcacag	cagctgcgag	agatcttcaa	catggcgagg	1380
gatcagcgca	gccgagccac	cttgatgctg	ctgagccacg	cggccaacaa	ctatcataag	1440
acaaacccta	cagggaccca	ggagctgctg	gacattgcta	attacctgat	ggaacagatt	1500
caagatgact	gcaactggga	tgaagattac	acctatttga	ttctgcgggt	cattggaaat	1560
atgggccaaa	ccatggagca	gttaactcca	gaactcaagt	cttcaatcct	gaaatgtgtc	1620
caaagtacaa	agccatcact	gatgatccag	aaagctgcca	tccaggctct	gcggaaaaatg	1680
gagcctaaag	acaaggacca	ggaggttctt	cttcagactt	tccttgatga	tgcttctccg	1740
ggagataagc	gactggctgc	ctatcttatg	ttgatgagga	gtccttcaca	ggcagatatt	1800
aacaaaattg	tccaaattct	accatgggaa	cagaatgagc	aagtgaagaa	ctttgtggct	1860
tcccatattg	ccaatatctt	gaactcagaa	gaattggata	tccaagatct	gaaaaagtta	1920
gtgaaagaag	ttctgaaaga	atctcaactt	ccaactgtca	tggaacttcag	aaaattctct	1980
cggaaactatc	aactctacaa	atctgtttct	attccatcac	ttgacccagc	ctcagccaaa	2040
atagaaggga	atcttatatt	tgatccaaat	aactaccttc	ctaaagaaag	catgctgaaa	2100
actaccctca	ctgcctttgg	atttgcttca	gctgacctca	tcgagattgg	cttggaagga	2160
aaaggctttg	agccaacatt	ggaagctctt	tttgggaagc	aaggattttt	cccagacagt	2220
gtcaacaaaag	ctttgtactg	ggttaatggt	caagttcctg	atggtgtctc	taaggtctta	2280
gtggaccact	ttggctatac	caaagatgat	aaacatgagc	aggatatggt	aaatggaata	2340
atgctcagtg	ttgagaagct	gattaaagat	ttgaaatcca	aagaagtccc	ggaagccaga	2400
gcctacctcc	gcatcttggg	agaggagctt	gggtttgcca	gtctccatga	cctccagctc	2460
ctgggaaaagc	tgcttctgat	gggtgcccgc	actctgcagg	ggatccccc	gatgattgga	2520
gaggtcatca	ggaagggctc	aaagaatgac	ttttttcttc	actacatctt	catggagaat	2580
gcctttgaac	tcctccactg	agctggatta	cagttgcaaa	tatcttcac	tggagtcatt	2640
gctcccggag	ccaaggctgg	agtaaaactg	gaagtagcca	acatgcaggc	tgaactgggtg	2700
gcaaaaccct	ccgtgtctgt	ggagtttgtg	acaaatatgg	gcatcatcat	tccggacttc	2760
gctaggagtg	gggtccagat	gaacaccaac	ttcttccacg	agtcgggtct	ggaggctcat	2820
gttgccctaa	aacctgggaa	gctgaagttt	atcattcctt	ccccaaagag	accagtcaag	2880
ctgctcagtg	gaggcaacac	attacatttg	gtctctacca	ccaaaacgga	ggtgatccca	2940
cctctcattg	agaacaggca	gtcctggtca	gtttgcaagc	aagtctttcc	tggcctgaat	3000
tactgcacct	caggcgctta	ctccaacgcc	agctccacag	actccgctc	ctactatccg	3060

ctgaccgggg	acaccagatt	agagctggaa	ctgaggccta	caggagagat	tgagcagtat	3120
tctgtcagcg	caacctatga	gtccagaga	gaggacagag	ccttggtgga	tacctgaag	3180
tttgtaactc	aagcagaagg	tgcaagcag	actgaggcta	ccatgacatt	caaatataat	3240
cggcagagta	tgacctgtgc	cagtgaagtc	caaattccgg	atthttgatgt	tgacctcgga	3300
acaatcctca	gagttaatga	tgaatctact	gagggcaaaa	cgtcttacag	actcaccctg	3360
gacattcaga	acaagaaaat	tactgaggtc	gccctcatgg	gccacctaag	ttgtgacaca	3420
aaggaagaaa	gaaaaatcaa	gggtgttatt	tccatacccc	gtttgcaagc	agaagccaga	3480
agtgagatcc	tcgcccactg	gtcgccctgcc	aaactgcttc	tccaaatgga	ctcatctgct	3540
acagcttatg	gtccacacagt	ttccaagagg	gtggcatggc	attatgatga	agagaagatt	3600
gaatttgaat	ggaacacagg	caccaatgta	gataccaaaa	aaatgacttc	caatttcctt	3660
gtggatctct	ccgattatcc	taagagcttg	catatgtatg	ctaatagact	cctggatcac	3720
agagtccttc	aaacagacat	gactttccgg	cacgtgggtt	ccaaattaat	agttgcaatg	3780
agctcatggc	ttcagaaggc	atctgggagt	cttccttata	cccagacttt	gcaagaccac	3840
ctcaatagcc	tgaaggagtt	caacctccag	aacatgggat	tgccagactt	ccacatccca	3900
gaaaacctct	tcttaaaaag	cgatggccgg	gtcaaatata	ccttgaacaa	gaacagtttg	3960
aaaattgaga	ttcctttgcc	ttttgggtggc	aaatcctcca	gagatctaaa	gatgttagag	4020
actgttagga	caccagccct	ccacttcaag	tctgtgggat	tccatctgcc	atctcgagag	4080
ttccaagtcc	ctacttttac	cattcccaag	ttgtatcaac	tgcaagtgcc	tctcctgggt	4140
gttctagacc	tctccacgaa	tgtctacagc	aacttgtaca	actggtccgc	ctcctacagt	4200
gggtggcaaca	ccagcacaga	ccatttcagc	cttcgggctc	gttaccacat	gaaggctgac	4260
tctgtggttg	acctgctttc	ctacaatgtg	caaggatctg	gagaaacaac	atatgaccac	4320
aagaatacgt	tcacactatc	atgtgatggg	tctctacgcc	acaaatttct	agattcgaat	4380
atcaaattca	gtcatgtaga	aaaacttgga	aacaaccag	tctcaaaagg	tttactaata	4440
ttcgatgcat	ctagtctctg	gggaccacag	atgtctgctt	cagttcattt	ggactccaaa	4500
aagaaacagc	atttgtttgt	caaagaagtc	aagattgatg	ggcagttcag	agtctcttcg	4560
ttctatgcta	aaggcacata	tggcctgtct	tgtcagaggg	atcctaacac	tggccggctc	4620
aatggagagt	ccaacctgag	gtttaactcc	tctcactctc	aaggcaccaa	ccagataaca	4680
ggaagatatg	aagatggaac	cctctccctc	acctccacct	ctgatctgca	aagtggcatc	4740
attaaaaata	ctgcttccct	aaagtatgag	aactacgagc	tgactttaaa	atctgacacc	4800
aatgggaagt	ataagaactt	tgccacttct	aacaagatgg	atatgacctt	ctctaagcaa	4860
aatgcactgc	tgcgttctga	atatcaggct	gattacgagt	cattgaggtt	cttcagcctg	4920
ctttctggat	cactaaattc	ccatgggtctt	gagttaaaatg	ctgacatctt	aggcactgac	4980
aaaattaata	gtgggtgctca	caaggcgaca	ctaaggattg	gccaagatgg	aatatctacc	5040
agtgaacga	ccaacttgaa	gtgtagtctc	ctgggtgctg	agaatgagct	gaatgcagag	5100
cttggcctct	ctggggcatc	tatgaaatta	acaacaaatg	gccgcttcag	ggaacacaat	5160
gcaaaattca	gtctggatgg	gaaagccgcc	ctcacagagc	tatcactggg	aagtgettat	5220
caggccatga	ttctgggtgt	cgacagcaaa	aacattttca	acttcaagggt	cagtcaagaa	5280
ggacttaagc	tctcaaatga	catgatgggc	tcatatgctg	aaatgaaatt	tgaccacaca	5340
aacagtctga	acattgcagg	cttatcactg	gacttctctt	caaaacttga	caacattttac	5400
agctctgaca	agttttataa	gcaaactgtt	aattttacagc	tacagcccta	ttctctggta	5460
actactttaa	acagtgaacct	gaaatacaat	gctctggatc	tcaccaacaa	tgggaaacta	5520
cggctagaac	ccctgaagct	gcatgtggct	ggtaacctaa	aaggagccta	ccaaaataat	5580
gaaataaaac	acatctatgc	catctcttct	gctgccttat	cagcaagcta	taaagcagac	5640
actgttgcta	aggttcaggg	tgtggagttt	agccatcggc	tcaacacaga	catcgctggg	5700
ctggcttcag	ccattgacat	gagcacaaac	tataattcag	actcactgca	tttcagcaat	5760
gtcttccggt	ctgtaatggc	cccgtttacc	atgaccatcg	atgcacatac	aatggcaat	5820
gggaaactcg	ctctctgggg	agaacatact	gggcagctgt	atagcaaatt	cctgttgaaa	5880
gcagaacctc	tggcattttac	tttctctcat	gattacaaag	gtccacaaag	tcatcatctc	5940
gtgtctagga	aaagcatcag	tgcagctctt	gaacacaaag	tcagtgccct	gcttactcca	6000
gctgagcaga	caggcacctg	gaaactcaag	acccaattta	acaacaatga	atacagccag	6060
gacttggatg	cttacaacac	taaagataaa	attggcgctg	agcttactgg	acgaactctg	6120
gctgacctaa	ctctactaga	ctccccatt	aaagtgccac	ttttactcag	tgagcccatc	6180
aatatcattg	atgctttaga	gatgagagat	gccgttgaga	agccccaaga	atttacaatt	6240
gttgcttttg	taaagtatga	taaaaaccaa	gatgttcaact	ccattaacct	cccatttttt	6300
gagaccttgc	aagaatattt	tgagaggaat	cgacaaaacca	ttatagttgt	actggaaaac	6360

gtacagagaa	acctgaagca	catcaatatt	gatcaatttg	taagaaaata	cagagcagcc	6420
ctgggaaaac	tcccacagca	agctaattgat	tatctgaatt	cattcaattg	ggagagacaa	6480
gtttcacatg	ccaaggagaa	actgactgct	ctcacaaaaa	agtatagaat	tacagaaaat	6540
gatatacaaa	ttgcattaga	tgatgccaaa	atcaacttta	atgaaaaact	atctcaactg	6600
cagacatata	tgatacaatt	tgatcagtat	attaaagata	gttatgattt	acatgatttg	6660
aaaatagcta	ttgctaatat	tattgatgaa	atcattgaaa	aattaaag	tcttgatgag	6720
cactatcata	tccgtgtaaa	tttagtaaaa	acaatccatg	atctacattt	gtttattgaa	6780
aatattgatt	ttaacaaaag	tggaagtagt	actgcaccc	ggattcaaaa	tgtggatact	6840
aagtaccaa	tcagaatcca	gatacaagaa	aaactgcagc	agcttaagag	acacatacag	6900
aatatagaca	tccagcacct	agctggaaaag	ttaaaacaac	acattgaggg	tattgatgtt	6960
agagtgtctt	tagatcaatt	gggaactaca	atttcatttg	aaagaataaa	tgatgttctt	7020
gagcatgtca	aacactttgt	tataaatctt	attggggatt	ttgaagtagc	tgagaaaatc	7080
aatgccttca	gagccaaagt	ccatgagtta	atcgagaggt	atgaagtaga	ccaacaaatc	7140
cagggtttta	tggtataaatt	agtagagttg	gccaccaat	acaagttgaa	ggagactatt	7200
cagaagctaa	gcaatgtcct	acaacaagtt	aagataaaaag	attactttga	gaaattgggt	7260
ggatttattg	atgatgctgt	caagaagctt	aatgaattat	cttttaaaac	attcattgaa	7320
gatgttaaca	aattccttga	catgttgata	aagaaattaa	agtcatttga	ttaccaccag	7380
tttgtatag	aaaccaatga	caaaatccgt	gaggtgactc	agagactcaa	tggtgaaatt	7440
caggctctgg	aactaccaca	aaaagctgaa	gcattaaaac	tggttttaga	ggaaaccaag	7500
gccacagttg	cagtgtatct	ggaaagccta	caggacacca	aaataacctt	aatcatcaat	7560
tggttacagg	aggctttaag	ttcagcatct	ttggctcaca	tgaaggccaa	attccgagag	7620
actctagaag	atacacgaga	ccgaatgtat	caaattggaca	ttcagcagga	acttcaacga	7680
tacctgtctc	tggtaggcca	ggtttatagc	acacttgctc	cctacatttc	tgattggtgg	7740
actcttgctg	ctaagaacct	tactgacttt	gcagagcaat	attctatcca	agattgggct	7800
aaacgtatga	aagcattggg	agagcaagg	ttcactgttc	ctgaaatcaa	gaccatcctt	7860
gggaccatgc	ctgcctttga	agtcagtcct	caggctcttc	agaaagctac	cttcagaca	7920
cctgatttta	tagtccccct	aacagatttg	aggattccat	cagttcagat	aaacttcaa	7980
gacttaaaaa	atataaaaa	cccatccagg	ttttccacac	cagaatttac	catccttaac	8040
accttccaca	ttccttcctt	tacaattgac	ttttagaaaa	tgaaagtaaa	gatcatcaga	8100
accattgacc	agatgctgaa	cagtgaagctg	cagtggcccg	ttccagatat	atatctcagg	8160
gatctgaagg	tgaggagacat	tcctctagcg	agaatcaccc	tgccagactt	ccgtttacca	8220
gaaatcgcaa	ttccagaatt	cataatccca	actctcaacc	ttaatgattt	tcaagttcct	8280
gaccttcaca	taccagaatt	ccagcttccc	cacatctcac	acacaattga	agtacctact	8340
tttggcaagc	tatacagtat	tctgaaaatc	caatctcctc	ttttcacatt	agatgcaaat	8400
gctgacatag	ggaatggaac	cacctcagca	aacgaagcag	gtatcgagc	ttccatcact	8460
gccaaaggag	agtccaaatt	agaagttctc	aattttgatt	ttcaagcaaa	tgcaaacctc	8520
tcaaacccta	agattaatcc	gctggctctg	aaggagtcag	tgaagttctc	cagcaagtac	8580
ctgagaacgg	agcatgggag	tgaaatgctg	ttttttggaa	atgctattga	gggaaaatca	8640
aacacagtg	caagtttaca	cacagaaaaa	aatacactgg	agcttagtaa	tggaagtatt	8700
gtcaagataa	acaatcagct	taccctggat	agcaacacta	aatacttcca	caaattgaac	8760
atccccaac	tggaacttctc	tagtcaggct	gacctgcgca	acgagatcaa	gacactgttg	8820
aaagctggcc	acatagcatg	gacttcttct	ggaaaagggt	catggaaatg	ggcctgcccc	8880
agattctcag	atgagggaac	acatgaatca	caaattagtt	tcaccataga	aggacccctc	8940
acttcccttg	gactgtccaa	taagatcaat	agcaaacacc	taagagtaaa	ccaaaacttg	9000
gtttatgaat	ctggctccct	caacttttct	aaacttgaaa	ttcaatcaca	agtcgattcc	9060
cagcatgtgg	gccacagtgt	tctaactgct	aaaggcatgg	cactgttttg	agaagggaag	9120
gcagagttta	ctggggaggca	tgatgctcat	ttaaatggaa	agggtatttg	aactttgaaa	9180
aattctcttt	tcttttcagc	ccagccattt	gagatcacgg	catccacaaa	caatgaagg	9240
aatttgaaag	ttcgttttcc	attaaggtta	acagggaaga	tagacttctc	gaataactat	9300
gcactgtttc	tgagtcccag	tgcccagcaa	gcaagttggc	aagtaagtgc	taggttcaat	9360
cagtataagt	acaacaaaaa	tttctctgct	ggaaacaacg	agaacattat	ggaggcccat	9420
gtaggaataa	atggagaagc	aatctggat	ttcttaaaaa	ttcctttaac	aattcctgaa	9480
atgctgtctac	cttacacaat	aatcacaaat	cctccactga	aagatttctc	tctatgggaa	9540
aaaacaggct	tgaagggaatt	cttgaaaaacg	acaaagcaat	catttgattt	aagtgtaaaa	9600
gctcagtata	agaaaaacaa	acacaggcat	tccatcacaa	atcctttggc	tgtgctttgt	9660

gagtttatca	gtcagagcat	caaateccttt	gacaggcatt	ttgaaaaaaa	cagaaacaat	9720
gcattagatt	ttgtcaccaa	atcctataat	gaaacaaaaa	ttaagtttga	taagtacaaa	9780
gctgaaaaat	ctcacgacga	gctccccagg	acctttcaaa	ttcctggata	cactgttcca	9840
gttgtaaatg	ttgaagtgtc	tccattcacc	atagagatgt	cggcattcgg	ctatgtgttc	9900
ccaaaagcag	tcagcatgcc	tagtttctcc	atcctagggt	ctgacgtccg	tgtgccttca	9960
tacacattaa	tcttgccatc	attagagctg	ccagtccttc	atgtccctag	aaatctcaag	10020
ctttctcttc	cagatttcaa	ggaattgtgt	accataagcc	atatttttat	tccctgccatg	10080
ggcaatatta	cctatgattt	ctcctttaaa	tcaagtgtca	tcacactgaa	taccaatgct	10140
gaacttttta	accagtcaga	tattgttgct	catctccttt	cttcatcttc	atctgtcatt	10200
gatgcactgc	agtacaaatt	agagggcacc	acaagattga	caagaaaaag	gggattgaag	10260
ttagccacag	ctctgtctct	gagcaacaaa	tttgtggagg	gtagtcataa	cagtactgtg	10320
agcttaacca	cgaaaaatat	ggaagtgtca	gtggcaacaa	ccacaaaagc	ccaaattcca	10380
attttgagaa	tgaatttcaa	gcaagaactt	aatggaaata	ccaagtcaaa	acctactgtc	10440
tcttctcca	tggaaattta	gtatgatttc	aattcttcaa	tgctgtactc	taccgctaaa	10500
ggagcagttg	accacaagct	tagcttgga	agcctcacct	cttacttttc	cattgagtca	10560
tctaccaaag	gagatgtcaa	gggttcgggt	ctttctcggg	aatattcagg	aactattgct	10620
agtgaggcca	acacttactt	gaattccaag	agcacacggg	cttcagtga	gctgcagggc	10680
acttccaaaa	ttgatgatat	ctggaacctt	gaagtaaaag	aaaattttgc	tggagaagcc	10740
acactccaac	gcatatatct	cctctgggag	cacagtacga	aaaaccactt	acagctagag	10800
ggcctctttt	tcaccaacgg	agaacatata	agcaaaagcca	ccctggaact	ctctccatgg	10860
caaatgtcag	ctcttgttca	ggtccatgca	agtcagccca	gttccttcca	tgatttccct	10920
gaccttggcc	aggaagtggc	cctgaatgct	aacactaaga	accagaagat	cagatggaaa	10980
aatgaagtcc	ggattcattc	tgggtctttc	cagagccagg	tcgagctttc	caatgaccaa	11040
gaaaaggcac	accttgacat	tgcaggatcc	ttagaaggac	acctaagggt	cctcaaaaat	11100
atcatcctac	cagtctatga	caagagctta	tgggatttcc	taaagctgga	tgtaaccacc	11160
agcattggta	ggagacagca	tcttcgtgtt	tcaactgcct	ttgtgtacac	caaaaacccc	11220
aatggctatt	cattctccat	ccctgtaaaa	gttttggtg	ataaattcat	tattcctggg	11280
ctgaaactaa	atgatctaaa	ttcagttctt	gtcatgccta	cgttccatgt	cccatttaca	11340
gatcttcagg	ttccatcgtg	caaacttgac	ttcagagaaa	tacaaatcta	taagaagctg	11400
agaacttcat	catttgccct	caacctacca	acactccccg	aggtaaaatt	ccctgaagtt	11460
gatgtgttaa	caaaaatttc	tcaaccagaa	gactccttga	ttcccttttt	tgagataacc	11520
gtgcctgaat	ctcagttaac	tgtgtcccag	ttcacgcttc	caaaaagtgt	ttcagatggc	11580
attgctgctt	tggatctaaa	tcagtagcc	aacaagatcg	cagactttga	gttggcccacc	11640
atcatcgtgc	ctgagcagac	cattgagatt	ccctccatta	agttctctgt	acctgctgga	11700
attgccattc	cttcctttca	agcactgact	gcacgctttg	aggtagactc	tcccgtgtat	11760
aatgccactt	ggagtgccag	tttgaaaaac	aaagcagatt	atgttgaaac	agtcctggat	11820
tccacatgca	gctcaaccgt	acagttccta	gaatatgaac	ttaatgtttt	gggaacacac	11880
aaaatcgaa	atggtacgtt	agcctctaag	actaaaggaa	catttgacac	ccgtgacttc	11940
agtgcagaat	atgaagaaga	tggcaaatat	gaaggacttc	aggaatggga	aggaaaagcg	12000
cacctcaata	tcaaaagccc	agcgttcacc	gatctccatc	tgcgctacca	gaaagacaag	12060
aaaggcatct	ccacctcagc	agcctcccca	gccgtaggca	ccgtgggcat	ggatatggag	12120
gaagatgacg	acttttctaa	atggaacttc	tactacagcc	ctcagtcctc	tccagataaa	12180
aaactcacca	tattcaaaac	tgagttgagg	gtccgggaat	ctgatgagga	aactcagatc	12240
aaagttaatt	gggaagaaga	ggcagcttct	ggcttgctaa	cctctctgaa	agacaacgtg	12300
ccaaggcca	caggggtcct	ttatgattat	gtcaacaagt	accactggga	acacacaggg	12360
ctcacctga	gagaagtgtc	ttcaaagctg	agaagaaatc	tgcaggacca	tgtgtgagtg	12420
gtttatcaag	gggccattag	ggaaattgat	gatatcgacg	agaggttcca	gaaaggagcc	12480
agtgggacca	ctgggacctc	ccaagagtgg	aaggacaagg	cccagaatct	gtaccaggaa	12540
ctggtgactc	aggaaggcca	agccagtttc	cagggactca	aggataacgt	gtttgatggc	12600
ttggtacgag	ttactcaaga	attccatatt	aaagtcaagc	atctgattga	ctcactcatt	12660
gattttctga	acttccccag	attccagttt	ccggggaaac	ctgggatata	cactaggagg	12720
gaactttgca	ctatgttcat	aagggaggta	gggacgggtac	tgtcccaggt	atattcgaaa	12780
gtccataatg	gttcagaaat	actgttttcc	tatttccaag	acctagtgtg	tacacttccct	12840
ttcgagttaa	ggaaacataa	actaatagat	gtaatctcga	tgtataggga	actgttgaaa	12900
gatttatcaa	aagaagccca	agaggtattt	aaagccattc	agtctctcaa	gaccacagag	12960

gtgctacgta	atcttcagga	ccttttacaa	ttcattttcc	aactaataga	agataacatt	13020
aaacagctga	aagagatgaa	atctacttat	cttattaatt	atatccaaga	tgagatcaac	13080
acaatcttca	atgattatat	cccatatggt	tttaaattgt	tgaaagaaaa	cctatgcctt	13140
aatcttcata	agttcaatga	atcttattcaa	aacgagcttc	aggaagcttc	tcaagagtta	13200
cagcagatcc	atcaatacat	tatggccctt	cgtgaagaat	atcttgatcc	aagtatagtt	13260
ggctggacag	tgaaatatta	tgaacttgaa	gaaaagatag	tcagtctgat	caagaacctg	13320
ttagttgctc	ttaaggactt	ccattctgaa	tatattgtca	gtgcctctaa	ctttacttcc	13380
caactctcaa	gtcaagttga	gcaatttctg	cacagaaata	ttcaggaata	tcttagcatc	13440
cttaccgatc	cagatggaaa	agggaaagag	aagattgcag	agctttctgc	cactgctcag	13500
gaaataatta	aaagccaggc	cattgcgacg	aagaaaataa	tttctgatta	ccaccagcag	13560
tttagatata	aactgcaaga	tttttcagac	caactctctg	attactatga	aaaatttatt	13620
gctgaatcca	aaagattgat	tgacctgtcc	attcaaaact	accacacatt	tctgatatac	13680
atcacggagt	tactgaaaaa	gctgcaatca	accacagtc	tgaaccctta	catgaagctt	13740
gctccaggag	aacttactat	catcctctaa	ttttttaaaa	gaaatcttca	tttattcttc	13800
ttttccaatt	gaactttcac	atagcacaga	aaaaattcaa	aatgcctata	ttgatcaaac	13860
catacagtga	gccagccttg	cagtaggcag	tagactataa	gcagaagcac	atatgaactg	13920
gacctgcacc	aaagctggca	ccagggctcg	gaaggctctc	gaactcagaa	ggatggcatt	13980
ttttgcaagt	taaagaaaat	caggatctga	gttattttgc	taaacttggg	ggaggaggaa	14040
caaataaatg	gagtccttat	tgtgtatcat				14070

<210> 3
 <211> 3805
 <212> DNA
 <213> Homo Sapiens

<220>
 <221> exon
 <222> (71)...(114)
 <223> Exon 1

<221> exon
 <222> (872)...(937)
 <223> Exon 2

<221> exon
 <222> (2031)...(2223)
 <223> Exon 3

<221> exon
 <222> (2805)...(3663)
 <223> Exon 4

<400> 3

cctatccctg	ggggaggggg	cgggacaggg	ggagccctat	aattggacaa	gtctgggac	60
cttgagtcc	actcagcccc	agcggagggtg	aaggacgtcc	ttccccagga	gccgggtgaga	120
agcgcagtcg	ggggcacggg	gatgagctca	ggggcctcta	gaaagagctg	ggaccctggg	180
aagccctggc	ctccaggtag	tctcaggaga	gtactcggg	gtcgggcttg	gggagaggag	240
gagcgggggt	gaggcaagca	gcaggggact	ggacctggga	agggctgggc	agcagagacg	300
acccgacccg	ctagaagggtg	gggtggggag	agcagctgga	ctgggatgta	agccatagca	360
ggactccacg	agttgtcact	atcattatcg	agcacctact	gggtgtcccc	agtgtcctca	420
gatctccata	actggggagc	caggggcagc	gacacggtag	ctagccgtcg	attggagaac	480
tttaaaatga	ggactgaatt	agctcataaa	tggaaacagg	cgcttaactg	tgaggttggg	540
gcttagaatg	tgaagggaga	atgaggaatg	cgagactggg	actgagatgg	aaccggcggt	600

ggggaggggg	tgggggggatg	gaatttgaac	cccgggagag	gaagatggaa	ttttctatgg	660
agggcgacct	ggggatgggg	agataagaga	agaccaggag	ggagttaa	agggaatggg	720
ttgggggagg	cttggtaa	gtgctgggat	taggctgttg	cagataatgc	aacaaggctt	780
ggaaggctaa	cctgggggtga	ggccgggttg	ggggcgctgg	gggtgggagg	agtcctcact	840
ggcgggtgat	tgacagtttc	tccttcccca	gactggccaa	tcacaggcag	gaagatgaag	900
gttctgtggg	ctgcgttgct	ggtcacattc	ctggcaggta	tggggcgggg	gcttgctcgg	960
ttccccccgc	tcctccccct	ctcatcctca	cctcaacctc	ctggccccat	tcagacagac	1020
cctggggccc	ctcttctgag	gcttctgtgc	tgttctctgg	ctctgaacag	cgatttgacg	1080
ctctctgggc	ctcggtttcc	cccatccttg	agataggagt	tagaagttgt	tttgttgttg	1140
ttgtttgttg	ttgtttgttt	gtttttttga	gatgaagtct	cgctctgtcg	cccaggctgg	1200
agtgcagtgg	cgggatctcg	gctcactgca	agctccgcct	cccagggtcca	cgccattctc	1260
ctgcctcagc	ctcccaagta	gctgggacta	caggcacatg	ccaccacacc	cgactaactt	1320
ttttgtat	tcagtagaga	cgggggtttca	ccatgttggc	caggctggtc	tggaaactcct	1380
gacctcaggt	gatctgccc	tttcgatctc	ccaaagtgtc	gggattacag	gcgtgagcca	1440
cgcacactgg	ctgggagtta	gaggtttcta	atgcattgca	ggcagatagt	gaataccaga	1500
cacggggcag	ctgtgatctt	tattctccat	cacccccaca	cagccctgcc	tggggcacac	1560
aaggacactc	aatacatgct	tttccgctgg	gccgggtggct	cacccctgta	atcccagcac	1620
tttgggaggc	caaggtggga	ggatcacttg	agcccaggag	ttcaacacca	gcctgggcaa	1680
catagtgaga	ccctgtctct	actaaaaata	caaaaattag	ccaggcatgg	tgccacacac	1740
ctgtgctctc	agctactcag	gaggctgagg	caggaggatc	gcttgagccc	agaagggtcaa	1800
ggttgacagt	aaccatgttc	agggcgctgc	actccagcct	gggtgacaga	gcaagaccct	1860
gtttataaat	acataatgct	ttccaagtga	ttaaaccgac	tccccctca	ccctgcccac	1920
catggctcca	aagaagcatt	tgtggagcac	cttctgtgtg	cccctaggta	gctagatgcc	1980
tggacggggg	cagaaggacc	ctgacccgac	cttgaacttg	ttccacacag	gatgccaggc	2040
caaggtggag	caagcgggtg	agacagagcc	ggagcccag	ctgcgccagc	agaccgagtg	2100
gcagagcgcc	cagcgtggg	aactggcact	gggtcgcttt	tgggattacc	tgcgctgggt	2160
gcagacactg	tctgagcagg	tgacaggagga	gctgtcagc	tcccaggtea	cccaggaaact	2220
gagggtgagt	tccccatcct	ggcccttgac	cctcctgggt	ggcggtata	cctccccagg	2280
tccagggttt	attctgcccc	tgtcgctaag	ctttgggggg	cctgggtctc	tgctgggtct	2340
agcttcctct	tcccatttct	gactcctggc	tttagctctc	tgggaattctc	tctctcagct	2400
ttgtctctct	ctcttccctt	ctgactcagt	ctctcacact	cgctcctggc	ctgtctctgt	2460
ccttccctag	ctcttttata	tagagacaga	gagatgggg	ctcactgtgt	tgcccaggct	2520
ggtcttgaac	ttctgggctc	aagcgatcct	ccgcctcgg	cctcccaaag	tgctgggatt	2580
agaggcatga	gcaccttgcc	cggcctccta	gctccttctt	cgtctctgcc	tctgcccctc	2640
gcactctgtc	tctgcactct	tctctgtctc	cttctctcgg	cctctgcccc	gttctctctc	2700
tccctcttgg	gtctctctgg	ctcatcccca	tctcgcccg	cccatcccag	ccctctctcc	2760
ccgcctcccc	actgtgcgac	accctcccg	cctctcggcc	gcagggcgct	gatggacgag	2820
accatgaagg	agttgaaggc	ctacaaatcg	gaactggagg	aacaactgac	ccgggtggcg	2880
gaggagacgc	gggcacggct	gtccaaggag	ctgcaggcgg	cgcaggcccg	gctgggcgcg	2940
gacatggagg	acgtgtgcgg	ccgcctgggt	cagtaccgcg	gcgaggtgca	ggccatgctc	3000
ggccagagca	ccgaggagct	gcgggtgcgc	ctcgccctcc	acctgcgcaa	gctgcgtaag	3060
cggctcctcc	gcgatgccga	tgacctgcag	aagcgctgg	cagtgtacca	ggccggggcc	3120
cgcgagggcg	ccgagcgcg	cctcagcgcc	atccgcgagc	gcctggggcc	cctgggtggaa	3180
cagggcccg	tgccggccgc	cactgtgggc	tccttgccg	gccagccgct	acaggagcgg	3240
gcccaggcct	ggggcgagcg	gctgcgcg	cggatggagg	agatgggcag	ccggaccgcg	3300
gaccgcctgg	acgaggtgaa	ggagcaggtg	gcggaggtgc	gcgccaagct	ggaggagcag	3360
gcccagcaga	tacgcctgca	ggccgaggcc	ttccaggccc	gcctcaagag	ctgggttcgag	3420
cccctgggtg	aagacatgca	gcgccagtgg	gccgggctgg	tggagaaggt	gcaggctgcc	3480
gtgggcacca	gcgccgcccc	tgtgcccagc	gacaatcact	gaacgccgaa	gcctgcagcc	3540
atgcgacccc	acgccacccc	gtgcctcctg	cctccgcgca	gcctgcagcg	ggagaccctg	3600
tccccgcccc	agccgtcctc	ctgggggtgga	ccctagttta	ataaagattc	accaagtttc	3660
acgcactctg	tggcctcccc	ctgtgatttc	ctctaagccc	cagcctcagt	ttctctttct	3720
gcccacatac	tgccacacaa	ttctcagccc	cctcctctcc	atctgtgtct	gtgtgtatct	3780
ttctctctgc	cctttttttt	ttttt				3805

<210> 4
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 4
agtctggatg ggtaagccgc cctca 25

<210> 5
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 5
ctgggctggc ttaagccatt gacat 25

<210> 6
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 6
gctctctggg gataacatac tgggc 25

<210> 7
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 7
gatgccgttg agtagcccca agaat 25

<210> 8
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 8
gagaggaatc gataaacat tatag 25

19

<210> 9
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 9
tgtaagaaaa taaagagcag ccctg 25

<210> 10
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 10
gcagccctgg gataactccc acagc 25

<210> 11
<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 11
gcaagctaatt gattagctga attcattcaa t 31

<210> 12
<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 12
caagttttcac atgcctagga gaaactgact g 31

<210> 13
<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 13
atatacaaat tgcattgagat gatgccaaaa t 31

<210> 14
<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 14
aaactatctc aactgtagac atatatgata c 31

<210> 15
<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 15
gctaataatta ttgattaaat cattgaaatt a 31

<210> 16
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 16
tgatgagcac tagcatatcc gtgta 25

<210> 17
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 17
ctgcagcagc tttagagaca catac 25

<210> 18
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 18
aacagtgagc ttagtggtgcc cgttc 25

21

<210> 19
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 19
cagacttccg ttaaccagaa atcgc

25

<210> 20
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 20
aaagggtcat ggtaatgggc ctgcc

25

<210> 21
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 21
acatatatga tataatttga tcagt

25

<210> 22
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 22
atggaggacg tgtgcggccg cctgg

25

<210> 23
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 23
gacctgcaga agtgcctggc agtgt

25

22

<210> 24
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Therapeutic Oligonucleotide Fragment

<400> 24
 gacctgcaga agcgctggc agtgt

25

<210> 25
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Therapeutic Oligonucleotide Fragment

<400> 25
 taaggtcagg agtttgagac cagcc

25

<210> 26
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Therapeutic Oligonucleotide Fragment-DNA and RNA

<400> 26
 agucuggaug ggtaagccgc ccuca

25

<210> 27
 <211> 39
 <212> DNA
 <213> R. Rattus

<400> 27
 aaagattcat gtgaaggaga tagtggggga ccccatgtt

39

<210> 28
 <211> 13
 <212> PRT
 <213> R. Rattus

<400> 28
 Lys Asp Ser Cys Glu Gly Asp Ser Gly Gly Pro His Val
 1 5 10

<210> 29
 <211> 39
 <212> DNA
 <213> R. Rattus

<400> 29
aaagattcat gtgaaggaga tcgtggggga ccccatgtt 39

<210> 30
<211> 68
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide-DNA and RNA

<400> 30
ggggtccccc acgatctcct tcacattttu gugaaggaga tcgtggggga ccccgcgcggt 60
tttcgcgc 68

<210> 31
<211> 68
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide-DNA and RNA

<400> 31
tgtgaaggag atcgtggggg accccttttg ggguccccca cgatcuccuu cacagcgcggt 60
tttcgcgc 68

<210> 32
<211> 68
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide-DNA and RNA

<400> 32
tgtgaaggag atcgtggggg accccttttg ggguccccca cgatcuccuu cacagcgcggt 60
tttcgcgc 68

<210> 33
<211> 68
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide-DNA and RNA

<400> 33
tgtcaaggag atcgtggggg accccttttg ggguccccca cgatcuccuu gacagcgcggt 60
tttcgcgc 68

<210> 34
<211> 25
<212> DNA
<213> Artificial Sequence

24

<220>
<223> PCR Primer

<400> 34
cattgctgac aaggaataca cgaac 25

<210> 35
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR primer

<400> 35
atttgccttt cattgcacac tcttc 25

<210> 36
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR primer

<400> 36
attgccttgc tggaactgga taac 24

<210> 37
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR primer

<400> 37
ttgcctttca ttgcacattc ttcac 25

<210> 38
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide Probe

<400> 38
aaggagatag tggggga 17

<210> 39
<211> 17
<212> DNA
<213> Artificial Sequence

25

<220>
<223> Oligonucleotide Probe

<400> 39
aaggagatcg tggggga 17

<210> 40
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide Probe

<400> 40
aaggagatag tggggga 17

<210> 41
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide Probe

<400> 41
aaggagatcg tggggga 17

<210> 42
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR primer

<400> 42
attgccttgc tggaactgga taaac 25

<210> 43
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR primer

<400> 43
ttgcctttca ttgcacattc ttcac 25

<210> 44
<211> 17
<212> DNA
<213> Artificial Sequence

26

<220>
 <223> Oligonucleotide Probe

 <400> 44
 aaggagatag tggggga 17

 <210> 45
 <211> 17
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Oligonucleotide Probe

 <400> 45
 aaggagatcg tggggga 17

 <210> 46
 <211> 23
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> PCR primer

 <400> 46
 gttgaccgag ccacatgcct tag 23

 <210> 47
 <211> 68
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Therapeutic Oligonucleotide-DNA and RNA

 <400> 47
 ctgggctggc ttaagccatt gacatuuua ugucaauggc uuaagccagc ccaggcgcg 60
 uuucgcgc 68

 <210> 48
 <211> 88
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Therapeutic Oligonucleotide-DNA and RNA

 <400> 48
 gacaagtttc acatgcctag gagaaactga ctgctuuua gcagucaguu ucucctaggc 60
 augugaaacu ugucgcgcgu uuucgcgc 88

 <210> 49
 <211> 76
 <212> DNA

27

<213> Artificial Sequence

<220>

<223> Therapeutic Oligonucleotide-DNA and RNA

<400> 49

ccctgaatgt cctggaaatg actgccgatt ttttaucttca gucattttcca ggacauucag 60
gggcgcggttt tcgcgc 76

<210> 50

<211> 80

<212> DNA

<213> Artificial Sequence

<220>

<223> Therapeutic Oligonucleotide-DNA and RNA

<400> 50

taagttttctt aactgggatt attagctggg gttttcccca gctaataatc ccagttaaga 60
aacuuagcgc gttttcgcgc 80

<210> 51

<211> 68

<212> DNA

<213> Artificial Sequence

<220>

<223> Therapeutic Oligonucleotide-DNA and RNA

<400> 51

gtttcttaac tgggattatt agctgttttc agcuaataat cccaguuaag aaacgcgcgt 60
tttcgcgc 68

<210> 52

<211> 80

<212> DNA

<213> Artificial Sequence

<220>

<223> Therapeutic Oligonucleotide-DNA and RNA

<400> 52

taagttttctt aactgggatt attagctggg guuuucccca gcuaauaaucc ccaguuaaga 60
aacuuagcgc guuuucgcgc 80

<210> 53

<211> 80

<212> DNA

<213> Artificial Sequence

<220>

<223> Therapeutic Oligonucleotide-DNA and RNA

<400> 53

28

taagtttctt aactgggatt attagctggg guuuucccca gcuaauaau ucaguuuaga 60
aacuuagcgc guuuucgcgc 80

<210> 54
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 54
ggcgaggaca tgtgcgaccg ggcgc 25

<210> 55
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 55
gagatgcgcg actgcgcgcg cgccc 25

<210> 56
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 56
cgcgaccgcg cgtgcgcgca tgtgg 25

<210> 57
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 57
agcgaccagc tgtgccagcg cttgg 25

<210> 58
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Therapeutic Oligonucleotide Fragment

<400> 58
gagctgcgcc agtgcttggc cgcgc

25

<210> 59
<211> 8966
<212> DNA
<213> Homo sapiens

<220>
<221> CDS
<222> (703)...(748)

<221> CDS
<222> (1678)...(2280)

<400> 59
ctaaagaaga gcactggtgg gaggacaggg cgggggaagg gggaggggag tgaagtagtc 60
tccttggaat gctggtggtg ggggaggcag tctccttggg ggaggagtc cagcgtccct 120
ccctccctc cctctgcca cacaatggac aatggcaact gcccacacac tcccatggag 180
gggaagggga tgagtgcagg gaaccccgac cccacccggg agacctgcaa gcctgcagca 240
ctccctccc gcccccactg aacccttgac cctgcccctg cagccccgc agcttgctgt 300
ttgcccactc ctatttgccc agtcccaggg acagagctga tccttgaact cttaagttcc 360
acattgccag gaccagtgg cagcaacagg gccagggctg ggcttatcag cctcccagcc 420
cagaccctgg ctgcagacat aaataggccc tgcaagagct ggctgcttag agactgcgag 480
aaggaggtgc gtccctgctgc ctgccccggc actctggctc cccagctcaa gggtcaggcc 540
ttgcccagg ccgggcctct ggggtacctga ggtcttctcc cgctctgtgc ccttctctc 600
acctggctgc aatgagtggt ggagcacggg gcttctgcat gctgaaggca cccactcag 660
ccaggccctt cttctcctcc aggtccccc cggccctca ggatgaaagc tggggtgctg 720
accttgccg tgctcttctc gacgggtagg tgtccctcaa cctaggagcc aaccatcggg 780
gggctttctc cctaaatccc cgtggcccac cctcctgggc agaggcagca gggttctcac 840
tgccccctc tccccacct ccaagcttgg ccttctggct cagatctcag cccacagctg 900
gcctgatctg ggtctccct cccaccctca gggagccagg ctccggcattt ctggcagcaa 960
gatgaacccc cccagagccc ctgggatcga gtgaaggacc tggccactgt gtacgtggat 1020
gtgctcaaag acagcggcag agactatgtg tcccagttt aaggctccgc cttgggaaaa 1080
cagctaaagt aaggacccag cctggggttg agggcagggg cagggggcag aggcctgtgg 1140
gatgatgttg aagccagact ggccgagtc tcacctaata tctgatgagc tgggccccac 1200
agatggtctg gatggagaaa ccggaatgga tctccaggca gggtcacagc ccatgtcccc 1260
tgcaaaggac agaccagggc tgcccgatgc gtgatcacag agccacattg tgccctgcaag 1320
tgtagcaagc ccctttccct tcttcaccac ctctctgtct cctgcccagc aagactgtgg 1380
gctgtcttcg gagaggagaa tgcgctggag gcatagaagc gaggtccttc aagggccac 1440
tttgagagac aacgtaactg ggcaccagtc ccagctctgt ctcttttta gctcctctct 1500
gtgcctcggc ccagctgcac aacggggcat ggcttggcgg ggcaggggtg ttggttgaga 1560
gtgtactgga aatgctaggc cactgcacct ccgcgacag gtgtcaccca gggctcacc 1620
ctgataggct ggggcgctgg gaggccagcc ctcaaccctt ctgtctcacc ctccagccta 1680
aagctccttg acaactggga cagcgtgacc tccaccttca gcaagctgcg cgaacagctc 1740
ggccctgtga cccaggagtt ctgggataac ctggaaaagg agacagaggg cctgaggcaa 1800
gagatgagca aggatctgga ggaggtgaag gccaaaggtg agccctacct ggacgacttc 1860
cagaagaagt ggcaggagga gatggagctc taccgccaga aggtggagcc gctgcgcgca 1920
gagctccaag agggcgcgcg ccagaagctg cacgagctgc aagagaagct gagccactg 1980
ggcgaggaga tgcgcgaccg cgcgcgcgcc catgtggacg cgctgcgcac gcatctggcc 2040
ccctacagcg acgagctgcg ccagcgttg gccgcgcgcc ttgaggctct caaggagaac 2100
ggcggcgcca gactggccga gtaccacgcc aaggccaccg agcatctgag cagctcagc 2160
gagaaggcca agcccgct cgaggacctc cgccaaggcc tgctgcccgt gctggagagc 2220
ttcaaggtca gcttctgag cgctctcgag gagtacacta agaagctcaa caccagtgga 2280
ggcgcccccc gccgcccccc ttcccggtgc tcagaataaa cgtttccaaa gtgggaagca 2340

gcttctttct	tttgggagaa	tagagggggg	tgcgggggaca	tccggggggag	cccggggaggg	2400
gccttttgcc	ctggagcagg	gacttcctgc	cggatctcaa	caactccgtg	cccagactgg	2460
acgtcttagg	gccaagatcg	acgttgagg	acctgctgga	cgcntggctg	cttacgagtg	2520
agggagtaga	gtctgcctta	gcaaggctca	agtagaaagg	aagtcacagc	ggacnaggca	2580
aagccacaga	caatccaagg	ccaggtgccc	tgaaggggg	tcaaacaagg	cctgcagcct	2640
gtctgaggcg	ggccaggaaa	cagcggttgc	tttagctggg	agcagtcggg	ttccccgtcc	2700
ccagaggtgt	gtccgtatag	agccttctcc	agcccagccg	ctgtcagcgg	gcgggacgga	2760
gcggggcgcc	tcaggagacc	agccactggg	attggggttt	gggtcccgggt	gcaagtgaag	2820
cgcttgaggt	ttgcgcctgt	cctcctttac	taattcaaaa	acctctcaaa	cagacacttc	2880
ccttttcttc	tcacaaggcc	agtatcccc	tcccactact	cccataccgc	ccagaaacag	2940
ccgcggcttc	ctcaggcaca	gcagtgggaag	ccagtcctcc	acccctcgcg	gctccatgcc	3000
atgccacccc	ctctttctgc	cagccctggc	agaagctggc	ctgagtaaga	aaattcacca	3060
ccacctcttg	caggtacatt	tttatttcca	agatgctctc	atatctgtgc	tctcactgca	3120
tcctcccttc	cccacatcct	ggctagattg	ccatcacagc	cagagcatgg	atgaggacac	3180
tgaagcctgg	acctgtgaag	tcgcttgccc	agtgaacagc	aggatgggct	aggcagcgct	3240
ttttagaccc	tgcacccctg	gccatccatg	attattgaaa	agagtgtgcg	gggtcgggtgc	3300
ggtggctcaa	gcctgtaatc	ccagcacttt	gggaggctga	ggtgggcgta	tcacttcagg	3360
gccaggagtt	tgagaccagc	ctggccaata	tggtgaaacc	ctgtctctac	taaaaatata	3420
aaaaaaaaat	cagctggcat	ggtggcttgc	accctgaatc	ccagctacta	ggaaggctga	3480
ggcaggagaa	tcgcttgaac	ctgggaggca	gaggtcacag	tgagccgaaa	tcatgccact	3540
gcactccagc	ctgggcgacg	gagcaagact	ccagctaaaa	aaaaaaaaaa	aaaaaaaaag	3600
agtgtgtggc	ctggcactca	agttcacatg	ggtgtgcagg	catgcctgtg	tattctcaca	3660
tgacctccct	gtcacgggtc	cctccttgca	ctcatgtctg	aatgtccccg	ctgcacgcac	3720
atggcttcac	agatctggca	gtgccttccc	tacctctctc	ctgcagggcc	ttttgcccc	3780
tcatgcaggc	ccctggataa	tcggccccc	ccccatgtcc	ccatctccag	tgtatcttag	3840
ctaccctagg	taaaggagtg	ggcttttttag	ttcctaacct	tccagagcta	caacagcagt	3900
catccagcca	ggtctgggtg	ggaacattht	ctagatacgg	gtgctgagat	ctctcagccc	3960
agagagaagc	cctgggggat	tttcagagag	aaagcagctc	ccaggtgggg	ctggatgtac	4020
tgatgccact	gagatctgta	aaggagtccc	taacacctga	cataggagtg	acaaaactgt	4080
tttctgcacc	aactgagcag	aatacacgca	gctgacctgg	gctcaagggtc	tggccctgcc	4140
acgtgctggc	tctgtgatgc	tggccaagtg	ccttcgcctc	tccggggccac	agttttttga	4200
tctgaagagt	ggagccctac	tcaagccatc	tgcagctctc	gggctctctg	acctgacatc	4260
tttcgggtgg	tggggacaca	aagggaagcag	cctctattgg	gagaccttgt	gcttcttttt	4320
ggtcccagga	cactgcccc	caccactcca	gtccgggtccc	aaggggccag	tcagctcaac	4380
tgtaatcatg	acaacattga	tcaagcatct	ttacgtgcag	gtgctgtgcc	aaacgggtcg	4440
aacgctctct	catttcaatc	tcacggcaaa	cctacgggtg	aggggggtacg	gttgtatcca	4500
ctttacatgt	aagaaactga	ggctgatatc	aagtgggtga	gccaagaata	gtgcctcggt	4560
gcactttact	ccaacctcta	gcccacccgg	cctcctccct	tcacgtgcgc	ctaagagggc	4620
taggtggcct	ggatagggga	ggtcagctcc	acagttttga	gtaaacacac	acagctctca	4680
ctctgatgac	acttaagtgc	caggcatagt	ggctggcatg	gggcacacac	tcaagtcatg	4740
ttgtgcagca	cctaacagtt	tatcaaagta	tcagcaaaact	tattgtcctg	tttgaccttc	4800
cgcacaaaagc	tgtcaaggaa	ggcaggggtac	ggaggggtgat	tcctacctta	gagatgaaga	4860
aactgaggcc	cagagactag	cccagctacc	agaagggtgga	tagagcgctg	gcctccatgc	4920
ctgcctgacc	tggagtctgt	ccagtgccca	cccacagaac	agcctcgccc	ctttcccatg	4980
cccagacaca	gatggcacac	ttgcgacggc	ccactcatag	cagcttcttg	tccagcttta	5040
ttgggaggcc	agcatgcctg	gagggggggc	aggcatgagg	tggggtagga	gagcactgag	5100
aatactgtcc	cttttaagca	acctacaggg	gaagccctgg	agattgcagg	acccaaggag	5160
ctggcaggat	ggataggcag	gtggacttgg	ggtattgagg	tctcaggcag	ccacggctga	5220
agttgggtctg	acctcaggg	ccaaatccca	gaactcagag	aacttgtcct	taacgggtgct	5280
ccagtagtct	ttcagggaac	tgaagccatc	ggtcacccag	cccctaaatc	agtcagggga	5340
agcaacagag	cagggcatga	gaactcctct	gtaggcaacc	atgggaccca	cacccatgtc	5400
cccactggac	gacaccagtc	aggaccacac	caccctctca	acttcactgg	acgacagccc	5460
tgagacctca	ggcagggaatc	ccccacagct	gccaccctgg	gagaagagat	atccttgcag	5520
gaaccccagc	acaagtcaaa	ccctgccatc	tccaggtcac	ctcagatgtt	tatgcccctg	5580
ggcctgaggc	acaaagtgac	aggggtggga	gatgtgaaag	gtcaaggctg	tcattgtttt	5640

ctgtgcaaac	agcaccacct	ggagttgcac	aacctgggtg	ctctgagcag	ggtaggacag	5700
agggaggcag	cctctcattt	ggaaagtcac	tggaggattg	attagttgtg	tgatctgggg	5760
caggtcacct	aatcgctctg	agcctcaatt	ttctcatctg	caaagtgaga	aaataacacc	5820
taccccaaag	cgggttctgg	ggactaagaa	tgtttatgaa	cacctctgct	atgccagcta	5880
atgccagtca	gggtgaggtg	gagaagggag	taggggagag	gagagtactg	atgggcaggg	5940
gcaggatggg	agaggaagga	cacacttctg	ctcacactgg	cccaggtctt	tagaacaaag	6000
cagaagcact	cacgggcttg	aattgggtca	ggtggggcct	cccaaggcaa	acccttagcc	6060
caggttcccc	aacccccaa	tcatagatag	gtactggtcc	atggactgtt	aggagccggg	6120
ctgcacagca	cgaagtgagt	ggtgggtgag	tgagcaatac	gcctgagctc	cgcctcctgt	6180
cagatcaagt	ggccttacat	tctcatagga	gtgtgaaccc	tattgtgaac	tgacatgcc	6240
agggatctag	gtggagggct	ccttatgaga	atctaattgc	tgatgatctg	aggtggaaca	6300
gcttcatccc	aaaaccatcc	ccgctggccc	atggaaaaat	tgtccaccac	aaaactggtc	6360
cctggtgcc	aaaaggctga	ggaccactgc	tctagccaga	ccttcagaaa	aggaaaatgg	6420
ggccaggcgc	agtggctcgt	gcctgtaatc	ccagcacttt	gggaggccga	ggcaggagga	6480
tccctgagg	tcaggaattc	aagaccaacc	tggccaacat	ggtgaaaccc	catctctgct	6540
aaaaatacaa	aaattagctg	ggtgtggtgg	cgcgtgcctg	taatcccagc	tacttgggag	6600
gctgaggcag	gagaatgggt	tgaaccgggg	agacggaggt	tgcaagtggc	cgagatggca	6660
ccactgcact	ccagcctggg	tgacagaggg	agactccatt	aaaaaaaaaa	gaagaagaaa	6720
gaaagagaga	aaggaagaga	gggaaagaaa	gaaaaaggaa	agaaagaaa	aaagagaaa	6780
aaagaaaaga	aagaaaagaaa	ggaaagaaa	aaaggaaa	aagaaaagaaa	gaaagaaa	6840
aagaaaagaaa	gaaagaaa	aagaaaata	gaaagaaa	gaaaggaaa	ggaaaggaaa	6900
atggaaatga	gggctaaaac	ggcgcgcccc	taggactgct	ccgggagaaa	ggagggacct	6960
gggaggaatg	cagtctcaac	tctgtcatct	ctcccgagc	agcctgacaa	aggccctgtg	7020
aggaaagagg	aggctgaaga	gcacaggccc	aggggaggct	ggagcacctc	cattccattg	7080
ttgggatctc	accagggcag	gggcgggtgg	gaatggaggc	agctggcagg	ggggatgggg	7140
agggaggcca	gcggtgttac	ctggcctgct	gggccacctg	ggactcctgc	acgctgctca	7200
gtgcatcctt	ggcggtcttg	gtggcggtgt	tcatgtagcc	ctgcatgaag	ctgagaaggg	7260
aggcatcctc	ggcctctgaa	gctcctgagg	aaagagcagg	gctgagtggg	gtggatcggc	7320
ctctggacga	gccctggggc	tcctgcttga	ccaccattg	gactgggatc	cccaagttgc	7380
ctccaccctg	ccccagccc	agtcccacca	agtgcctacg	ggcagaggcc	aggagcgcca	7440
ggagggcaac	aacaaggagt	acccggggct	gcatggcacc	tctgttctct	caagggaagt	7500
tctgtgagg	ggcaccacag	gtcccatg	ctctggaccc	ctccctgggg	aggtggcggt	7560
gcccctaagg	tagaacctta	gctgggtctg	ccagaaggag	taggggccc	ctcctgctca	7620
tacgggtctt	cagaaggggg	actggtgagc	ggcgagggat	cgaggcccaa	aggaggtggg	7680
tgggatggag	cagaaaaccc	accagactga	acatcaaggc	actgcggtct	ggactgatct	7740
ccgtccagtc	cagccaacat	gctgtgtgtc	tttgggtgat	ttctggccct	cttcaggcc	7800
tcagtttccc	tgtctgggg	aggactgggc	tgtctaaggc	ttcctccagc	cctaagcctg	7860
agaatgagg	ggggaacctg	cacttgagc	cacttccagc	cccacccctt	atgtagcttt	7920
gggcaagtga	cacctcccg	gcctccatgt	tcttcaggtt	atgatgagg	gtggggggca	7980
cccgctccagc	tccgaggctt	ccttagctct	agcaagtgtc	tctccaggct	tgctggctgg	8040
gctgggcagg	gagctcctct	tgccctctt	catcctcctc	ccctcctctt	tccctcccc	8100
agagggcatt	acctggagca	gctgcctcta	gggatgaact	gagcagacag	gcaggagggt	8160
tctgacctgt	tttatatcat	ctccagggca	gcaggcactg	aggaccag	gcgctgggca	8220
aaggtcacct	gctgaccagt	ggagatgagg	gcctgaggca	gggtgtccag	atgcagcaag	8280
cgggcgggag	agttgggaaa	tccctaggag	actgagtcca	cgctgctgtc	ccgccagccc	8340
tgacgcccag	atgagctcag	gaactggggg	tgggcctggg	gagcctcatt	ccttcctagc	8400
tgactggctc	cccagggaga	ggctggtgag	aggggaaatg	gcttggacat	aggccagcgc	8460
tcgggccc	ctcagccttt	cacactggaa	tttcaggccc	ctccctccac	cagccccaag	8520
ccctgaacac	agcctggagt	agaggggtga	ggggcttctt	cagacttgag	aacaagtggg	8580
tggcttgggc	tgggggggtg	ttggagtaaa	ggcacagaag	accgggcac	agtggctcgc	8640
aaggcctggt	tacagggtg	agctcacagc	ccctcccagc	acctccatct	ctgggtttca	8700
atccaggcag	gcgagtgtg	ctcaccagc	agccccccac	ccgccccccac	cctgtgtgcc	8760
ccaccgccgc	ctccccctga	gtgtagggca	ggggttgggt	gagaagcgca	aggccgctca	8820
gagcccgagg	cctttgcccc	tccctccacc	aggctcccta	ttttgccctc	tggacccact	8880
gataacatcc	ctgggggagg	agccccacc	aggactgatt	ctctcggtta	ccttcctggg	8940

ttcctcagtg gggctggggg aagagc

8966

INTERNATIONAL SEARCH REPORT

national application No.
PCT/US98/17908

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A01N 43/04; C07H 21/04; C12N 15/09, 15/11, 15/85

US CL :435/440, 455, 471, 490; 514/44; 536/23.1, 23.5, 24.1, 24.2

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/440, 455, 471, 490; 514/44; 536/23.1, 23.5, 24.1, 24.2

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, BIOSIS, EMBASE, MEDLINE, DERWENT

search terms: apo? ldl? low density lipoprotein? cholesterol? atheroscleros? clathrin?

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A,P	US 5,736,157 A (WILLIAMS) 07 April 1998, see entire document.	1-4, 7-9, 24-50
A,E	US 4,772,549 A (FROSSARD) 20 September 1988, see entire document.	1-4, 7-9, 24-50



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

22 DECEMBER 1998

Date of mailing of the international search report

05 FEB 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

NANCY J. DEGEN

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/17908

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 5-6, 10-23, 51-52
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest☐
☐

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.